



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



SMC6 (m): 293T Lysate: sc-123650

BACKGROUND

Breaks in double stranded DNA often arise during DNA replication or as a result of exposure to DNA-damaging agents. Quick and accurate repair of these breaks is crucial for cell survival and genomic stability. Structural maintenance of chromosomes (SMC) family members form heterodimeric complexes that modulate sister chromatid cohesion and chromosome condensation during mitosis. Two distinct SMC protein complexes are the SMC1/SMC3 heterodimer and the SMC2/SMC4 heterodimer. SMC5 and SMC6 play a crucial role in DNA repair as they form a complex that along with SUMO ligase, is also important in preventing DNA damage-induced apoptosis. This complex made up of SMC5 and SMC6 is crucial for sister chromatid homologous recombination DNA repair and also for prevention of chromosomal rearrangements.

REFERENCES

1. Lehmann, A.R. 2005. The role of SMC proteins in the responses to DNA damage. *DNA Repair* 4: 309-314.
2. Lee, K.M. and O'Connell, M.J. 2005. A new SUMO ligase in the DNA damage response. *DNA Repair* 5: 138-141.
3. Potts, P.R. and Yu, H. 2005. Human MMS21/NSE2 is a SUMO ligase required for DNA repair. *Mol. Cell. Biol.* 25: 7021-7032.
4. Watanabe, Y. 2005. The importance of being SMC5/6. *Nat. Cell Biol.* 7: 329-331.
5. Eydmann, T., Machin, F., Dalgaard, J.Z., Farmer, S., Jarmuz, A., Aragon, L. and Torres-Rosell, J. 2005. SMC5 and SMC6 genes are required for the segregation of repetitive chromosome regions. *Nat. Cell Biol.* 7: 412-419.
6. De Piccoli, G., Cortes-Ledesma, F., Ira, G., Torres-Rosell, J., Ceschia, A., Uhle, S., Farmer, S., Hwang, J.Y., Machin, F., McAleenan, A., Papusha, A., Cordon-Preciado, V., Clemente-Blanco, A., Vilella-Mitjana, F., Myung, K., Ullal, P., Jarmuz, A., Leitao, B., Bressan, D., Dotiwala, F., Zhao, X., Haber, J.E., Aguilera, A. and Aragon, L. 2006. SMC5-SMC6 mediate DNA double-strand-break repair by promoting sister-chromatid recombination. *Nat. Cell Biol.* 8: 1032-1034.
7. Lindroos, H.B., Ström, L., Itoh, T., Katou, Y., Shirahige, K. and Sjögren, C. 2006. Chromosomal association of the SMC5/6 complex reveals that it functions in differently regulated pathways. *Mol. Cell* 22: 755-767.
8. Potts, P.R., Porteus, M.H. and Yu, H. 2006. Human SMC5/6 complex promotes sister chromatid homologous recombination by recruiting the SMC1/3 cohesin complex to double-strand breaks. *EMBO J.* 25: 3377-3388.

CHROMOSOMAL LOCATION

Genetic locus: Smc6 (mouse) mapping to 12 A1.1.

PRODUCT

SMC6 (m): 293T Lysate represents a lysate of mouse SMC6 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

SMC6 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive SMC6 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.