



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Synaptojanin 2 (m): 293T Lysate: sc-123864

## BACKGROUND

The inositol polyphosphate 5-phosphatases selectively remove the phosphate from the 5-position of various phosphatidylinositols, which generate second messengers in response to extracellular signals. Synaptojanins are characterized by an N-terminal SAC1-like sequence, a central 5-phosphate domain and a unique C-terminal sequence and have been shown to use phosphatidylinositol 4,5-bisphosphate as a substrate. Synaptojanins exist as two isoforms, Synaptojanin 1 and 2, which differ in the C-terminal domain, and each isoform has multiple variants produced by alternative splicing. Synaptojanin 1 is expressed as two major forms of 145 kDa, which is found in brain, and 170 kDa, which is expressed in peripheral tissues. Eight splice variants of Synaptojanin 2 have been detected, including a brain specific isoform. Synaptojanins are thought to participate in the endocytosis of synaptic vesicles and the regulation of the actin cytoskeleton.

## REFERENCES

- Mitchell, C.A., et al. 1996. Regulation of second messengers by the inositol polyphosphate 5-phosphatases. *Biochem. Soc. Trans.* 24: 994-1000.
- Nemoto, Y., et al. 1997. Synaptojanin 2, a novel synaptojanin isoform with a distinct targeting domain and expression pattern. *J. Biol. Chem.* 272: 30817-30821.
- Zhang, X. and Majerus, P.W. 1998. Phosphatidylinositol signalling reactions. *Semin. Cell Dev. Biol.* 9: 153-160.
- Erneux, C., et al. 1998. The diversity and possible functions of the inositol polyphosphate 5-phosphatases. *Biochim. Biophys. Acta* 1436: 185-199.
- Khvotchev, M. and Sudhof, T.C. 1998. Developmentally regulated alternative splicing in a novel Synaptojanin. *J. Biol. Chem.* 273: 2306-2311.
- Seet, L.F., et al. 1998. Molecular cloning of multiple isoforms of Synaptojanin 2 and assignment of the gene to mouse chromosome 17A2-3.1. *Biochem. Biophys. Res. Commun.* 247: 116-22.
- Takenawa, T., et al. 1999. Regulation of phosphatidylinositol 4,5-bisphosphate levels and its roles in cytoskeletal re-organization and malignant transformation. *Chem. Phys. Lipids* 98: 13-22.
- Haffner, C., et al. 2000. Direct interaction of the 170 kDa isoform of Synaptojanin 1 with Clathrin and with the Clathrin adaptor AP-2. *Curr. Biol.* 10: 471-474.

## CHROMOSOMAL LOCATION

Genetic locus: Synj2 (mouse) mapping to 17 A1.

## PRODUCT

Synaptojanin 2 (m): 293T Lysate represents a lysate of mouse Synaptojanin 2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

Synaptojanin 2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Synaptojanin 2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Synaptojanin 2 (E-1): sc-390354 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Synaptojanin 2 expression in Synaptojanin 2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Synaptojanin 2 (E-1): sc-390354. Western blot analysis of Synaptojanin 2 expression in non-transfected: sc-117752 (A) and mouse Synaptojanin 2 transfected: sc-123864 (B) 293T whole cell lysates.



Synaptojanin 2 (D-11): sc-390247. Western blot analysis of Synaptojanin 2 expression in non-transfected: sc-117752 (A) and mouse Synaptojanin 2 transfected: sc-123864 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.