



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TACC3 (m2): 293T Lysate: sc-123891

BACKGROUND

TACC1 (transforming acidic coiled coil gene 1) is one of three TACC family members, which are thought to be involved in breast tumorigenesis. TACC1 is located on 8p11, in a chromosomal region that is amplified in approximately 15% of all breast tumor samples. The short arm of chromosome 8 also contains FGFR1, whose expression is enhanced in most breast cancer tumors. TACC family members TACC1, TACC2 and TACC3 map very closely to the corresponding FGFR1, FGFR2, FGFR3 genes on chromosomes 4, 8 and 10. Subsequently, since they are phylogenetically related, it is proposed that TACC and FGFR have similar roles in cell growth and differentiation. Also, TACC1 contains a conserved C-terminal region as in the *Drosophila* homolog D-TACC. D-TACC has been shown to be necessary for normal spindle function, and the mammalian TACC proteins appears to interact with centrosomes and microtubules in a similar manner.

REFERENCES

1. Dib, A., Adelaide, J., Chaffanet, M., Imbert, A., Le Paslier, D., Jacquemier, J., Gaudray, P., Theillet, C., Birnbaum, D. and Pebusque, M.J. 1995. Characterization of the region of the short arm of chromosome 8 amplified in breast carcinoma. *Oncogene* 10: 995-1001.
2. Yoshimura, N., Sano, H., Hashiramoto, A., Yamada, R., Nakajima, H., Kondo, M. and Oka, T. 1998. The expression and localization of fibroblast growth factor-1 (FGF-1) and FGF receptor-1 (FGFR-1) in human breast cancer. *Clin. Immunol. Immunopathol.* 89: 28-34.
3. Ugolini, F., Adelaide, J., Charafe-Jauffret, E., Nguyen, C., Jacquemier, J., Jordon, B., Birnbaum, D. and Pebusque, M.J. 1999. Differential expression assay of chromosome arm 8p genes identifies frizzled-related (FRP1/FRZB) and fibroblast growth factor receptor 1 (FGFR1) as candidate breast cancer genes. *Oncogene* 18: 1903-1910.
4. Still, I.H., Hamilton, M., Vince, P., Wolfman, A. and Cowell, J.K. 1999. Cloning of TACC1, an embryonically expressed, potentially transforming coiled coil containing gene, from the 8p11 breast cancer amplicon. *Oncogene* 18: 4032-4038.
5. Still, I.H., Vince, P. and Cowell, J.K. 1999. The third member of the transforming acidic coiled coil-containing gene family, TACC3, maps in 4p16, close to translocation breakpoints in multiple myeloma, and is upregulated in various cancer cell lines. *Genomics* 58: 165-170.

CHROMOSOMAL LOCATION

Genetic locus: Tacc3 (mouse) mapping to 5 B2.

PRODUCT

TACC3 (m): 293T Lysate represents a lysate of mouse TACC3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

TACC3 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TACC3 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.