



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TFIID (m2): 293T Lysate: sc-124002

BACKGROUND

In eukaryotic systems, initiation of transcription from protein-coding genes is a complex process requiring RNA polymerase II and broad families of auxiliary transcription factors. Such factors can be divided into two major functional classes: the basal factors that are required for transcription of all Pol II genes, including TFIIA, TFIIB, TFIID, TFIIE, TFIIIF and TFIIH; and sequence-specific factors that regulate gene expression. The basal transcription factors and Pol II form a specific multiprotein complex near the transcription start site by interacting with core promoter elements such as the TATA box generally located 25-30 base pairs upstream of the transcription start site. Binding of TFIID to the TATA element initiates assembly of the other factors into a pre-initiation complex. The TATA-binding subunit of TFIID (designated TFIID τ or TBP) from higher eukaryotes contains a highly conserved 180 amino acid C-terminal domain.

REFERENCES

- Maldonado, E., et al. 1990. Factors involved in specific transcription by mammalian RNA polymerase II: role of transcription factors IIA, IID, and IIB during formation of a transcription-competent complex. *Mol. Cell. Biol.* 10: 6335-6347.
- Peterson, M.G., et al. 1991. Structure and functional properties of human general transcription factor IIE. *Nature* 354: 369-373.
- Lee, D.K., et al. 1992. TFIIA induces conformational changes in TFIID via interactions with the basic repeat. *Mol. Cell. Biol.* 12: 5189-5196.
- Takada, R., et al. 1992. Identification of human TFIID components and direct interaction between a 250-kDa polypeptide and the TATA box-binding protein (TFIID τ). *Proc. Natl. Acad. Sci. USA* 89: 11809-11813.
- Huisinga, K.L., et al. 2007. A TATA binding protein regulatory network that governs transcription complex assembly. *Genome Biol.* 8: R46.
- Romier, C., et al. 2007. Crystal structure, biochemical and genetic characterization of yeast and *E. cuniculi* TAF₅ N-terminal domain: implications for TFIID assembly. *J. Mol. Biol.* 368: 1292-1306.
- Demeny, M.A., et al. 2007. Identification of a small TAF complex and its role in the assembly of TAF-containing complexes. *PLoS ONE* 2: E316.
- Bhattacharya, S., et al. 2007. Structural analysis and dimerization potential of the human TAF₅ subunit of TFIID. *Proc. Natl. Acad. Sci. USA* 104: 1189-1194.
- Holloway, K., et al. 2007. Action of transcription factors in the control of transferrin receptor expression in human brain endothelium. *J. Mol. Biol.* 365: 1271-1284.

CHROMOSOMAL LOCATION

Genetic locus: Tbp (mouse) mapping to 17 A2.

PRODUCT

TFIID (m2): 293T Lysate represents a lysate of mouse TFIID transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

TFIID (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TFIID antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

TFIID (C-8): sc-374035 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse TFIID expression in TFIID transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



TFIID (C-8): sc-374035. Western blot analysis of TFIID expression in non-transfected: sc-117752 (A) and mouse TFIID transfected: sc-124002 (B) 293T whole cell lysates.



TFIID (C-7): sc-271146. Western blot analysis of TFIID expression in non-transfected: sc-117752 (A) and mouse TFIID transfected: sc-124002 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.