



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TFIIH p34 (m): 293T Lysate: sc-124006

BACKGROUND

Initiation of transcription from protein-coding genes in eukaryotes is a complex process that requires RNA polymerase II, as well as families of basal transcription factors. Binding of the factor TFIID (TBP) to the TATA box is believed to be the first step in the formation of a multiprotein complex containing several additional factors, including TFIIA, TFIIB, TFIIE, TFIIIF and TFIIH. TFIIH (or BTF2) is a multisubunit transcription/DNA repair factor that possesses several enzymatic activities. The core of TFIIH is composed of five subunits, designated p89 (XPB or ERCC3), p62, p52, p44 and p34. Additional subunits of the TFIIH complex are p80 (XPD or ERCC2) and the ternary kinase complex composed of Cdk7, cyclin H and MAT1. Both p89 and p80 have ATP-dependent helicase activity. The p62, p52 and p44 subunits have been shown to be involved in nucleotide excision repair.

REFERENCES

1. Conaway, R.C., et al. 1989. An RNA polymerase II transcription factor has an associated DNA-dependent ATPase (dATPase) activity strongly stimulated by the TATA region of promoters. *Proc. Natl. Acad. Sci. USA* 86: 7356-7360.
2. Weeda, G., et al. 1990. A presumed DNA helicase encoded by ERCC3 is involved in the human repair disorders xeroderma pigmentosum and Cockayne's syndrome. *Cell*. 62: 777-791.
3. Weber, C.A., et al. 1990. ERCC2: cDNA cloning and molecular characterization of a human nucleotide excision repair gene with high homology to yeast Rad3. *EMBO J.* 9: 1437-1447.
4. Fischer, L., et al. 1991. Cloning of the 62-kilodalton component of basic transcription factor BTF2. *Science* 257: 1392-1395.
5. Gerard, M., et al. 1991. Purification and interaction properties of the human polymerase B (II) general transcription factor BTF2. *J. Biol. Chem.* 266: 20940-20945.
6. Flores, O., et al. 1992. Factors involved in specific transcription by mammalian RNA polymerase II. *J. Biol. Chem.* 267: 2786-2793.
7. Humbert, S., et al. 1994. p44 and p34 subunits of the STF2/TFIIH transcription factor have homologies with SSL1, a yeast protein involved in DNA repair. *EMBO J.* 13: 2393-2398.
8. Marinoni, J.C., et al. 1997. Cloning and characterization of p52, the fifth subunit of the core of the transcription/DNA repair factor TFIIH. *EMBO J.* 16: 1093-1102.

CHROMOSOMAL LOCATION

Genetic locus: Gtf2h3 (mouse) mapping to 5 F.

PRODUCT

TFIIH p34 (m): 293T Lysate represents a lysate of mouse TFIIH p34 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

TFIIH p34 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TFIIH p34 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.