



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



TFR2 (m): 293T Lysate: sc-124014

BACKGROUND

Iron is a vital molecule for living organisms because it is involved in a wide variety of metabolic processes, such as oxygen transport, DNA synthesis and electron transport. Excessive iron uptake leads to tissue damage as a result of formation of free radicals. Iron uptake and storage is tightly regulated by the feedback system of iron responsive element-containing gene products and iron regulatory proteins that modulate the expression levels of the genes involved in iron metabolism. The transferrin receptor 2 (TFR2) mediates the uptake of transferrin-bound iron. It is involved in iron metabolism, hepatocyte function and erythrocyte differentiation, and is highly expressed as a protein in liver as well as in hepatocytes and erythroid precursors. The gene encoding human TRF2 maps to chromosome 7q22.1 and is expressed as an α isoform, which encodes a transmembrane protein, and a β isoform, which encodes a shorter, intracellular protein. Mutations in the TFR2 gene result in hereditary hemochromatosis type III (HFE3), an iron overloading disorder that results in clinical complications, including cirrhosis, cardiopathy, diabetes, endocrine dysfunctions, arthropathy and susceptibility to liver cancer.

REFERENCES

1. Lieu, P.T., et al. 2001. The roles of iron in health and disease. *Mol. Aspects Med.* 22: 1-87.
2. Roetto, A., et al. 2001. New mutations inactivating transferrin receptor 2 in hemochromatosis type 3. *Blood* 97: 2555-2560.
3. Kawabata, H., et al. 2001. Regulation of expression of murine transferrin receptor 2. *Blood* 98: 1949-1954.
4. Kawabata, H., et al. 2001. Expression of transferrin receptor 2 in normal and neoplastic hematopoietic cells. *Blood* 98: 2714-2719.
5. Deaglio, S., et al. 2002. Structural, functional, and tissue distribution analysis of human transferrin receptor-2 by murine monoclonal antibodies and a polyclonal antiserum. *Blood* 100: 3782-3789.
6. Camaschella, C., et al. 2002. Genetic haemochromatosis: genes and mutations associated with iron loading. *Best Pract. Res. Clin. Haematol.* 15: 261-276.
7. Fletcher, L.M., et al. 2002. Haemochromatosis: understanding the mechanism of disease and implications for diagnosis and patient management following the recent cloning of novel genes involved in iron metabolism. *J. Intern. Med.* 251: 181-192.

CHROMOSOMAL LOCATION

Genetic locus: Trfr2 (mouse) mapping to 5 G2.

PRODUCT

TFR2 (m): 293T Lysate represents a lysate of mouse TFR2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

TFR2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TFR2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.