



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TGIF (m): 293T Lysate: sc-124025

## BACKGROUND

TGIF (for 5'-TG-3' interacting factor) was originally identified as a homeodomain protein that binds to a retinoid X receptor (RXR) responsive element, thereby inhibiting the binding of RXR to this site and repressing RXR-dependent transcriptional activation. TGIF is a member of the TALE (three amino acid loop extension) family of homeodomain-containing proteins. TGIF also binds to Smad2, to repress Smad2-Smad4-mediated transcription. Smad2, after phosphorylation mediated by TGF $\beta$  receptor, forms a complex with Smad4 and enters the nucleus to regulate transcription. The Smad2-Smad4 complex can interact with coactivators to form a transcriptional activation complex. Alternatively, the Smad2-Smad4 complex can interact with TGIF and HDACs to form a transcriptional repressor complex. Upon interaction with Smad2, TGIF is recruited to TGF $\beta$ -responsive genes, where it acts to repress TGF $\beta$ -induced transcription. The gene which encodes TGIF maps to human chromosome 18p11.3.

## REFERENCES

1. Bertolino, E., et al. 1995. A novel homeobox protein which recognizes a TGT core and functionally interferes with a retinoid-responsive motif. *J. Biol. Chem.* 270: 31178-31188.
2. Baker, J.C., et al. 1996. A novel mesoderm inducer, Madr2, functions in the activin signal transduction pathway. *Genes Dev.* 10: 1880-1889.
3. Janknecht, R., et al. 1996. TGF $\beta$ -stimulated cooperation of Smad proteins with the coactivators CBP/p300. *Genes Dev.* 12: 2114-2119.
4. Lana, G., et al. 1996. Partnership between DPC4 and Smad proteins in TGF $\beta$  signalling pathways. *Nature* 383: 832-836.
5. Burglin, T.R., et al. 1997. Analysis of TALE superclass homeobox genes (MEIS, PBC, KNOX, Iroquois, TGIF) reveals a novel domain conserved between plants and animals. *Nucleic Acids Res.* 25:4173-4180.
6. Poupnot, C., et al. 1998. Physical and functional interaction of Smads and p300/CBP. *J. Biol. Chem.* 273: 22865-22868.
7. Wotton, D., et al. 1999. A Smad transcriptional corepressor. *Cell* 97: 29-39.
8. Gripp, K.W., et al. 2000. Mutations in TGIF cause holoprosencephaly and link NODAL signalling to human neural axis determination. *Nat. Genet.* 25: 205-208.

## CHROMOSOMAL LOCATION

Genetic locus: Tgif1 (mouse) mapping to 17 E1.3.

## PRODUCT

TGIF (m): 293T Lysate represents a lysate of mouse TGIF transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## APPLICATIONS

TGIF (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TGIF antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20 $^{\circ}$  C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.