



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TNF-R1 (m): 293T Lysate: sc-124202

BACKGROUND

Tumor necrosis factor (TNF) is a pleiotropic cytokine whose function is mediated through two distinct cell surface receptors. These receptors, designated TNF-R1 and TNF-R2, are expressed on most cell types. The majority of TNF functions are primarily mediated through TNF-R1, while signaling through TNF-R2 occurs less extensively and is confined to cells of the immune system. Both of these proteins belong to the growing TNF and nerve growth factor (NGF) receptor superfamily, which includes FAS, CD30, CD27 and CD40. The members of this superfamily are type I membrane proteins that share sequence homology confined to the extracellular region. TNF-R1 shares a motif coined the "death domain" with FAS and three structurally unrelated signaling proteins, TRADD, FADD and RIP. This "death domain" is required for transduction of the apoptotic signal.

REFERENCES

- Derré, J., Kemper, O., Cherif, D., Nophar, Y., Berger, R. and Wallach, D. 1991. The gene for the type 1 tumor necrosis factor receptor (TNF-R1) is localized on band 12p13. *Hum. Genet.* 87: 231-233.
- Milatovich, A., Song, K., Heller, R.A. and Francke, U. 1991. Tumor necrosis factor receptor genes, TNFR1 and TNFR2, on human chromosomes 12 and 1. *Somat. Cell Mol. Genet.* 17: 519-523.
- Vielhauer, V., Stavakis, G. and Mayadas, T.N. 2005. Renal cell-expressed TNF receptor 2, not receptor 1, is essential for the development of glomerulonephritis. *J. Clin. Invest.* 115: 1199-1209.
- Dieudé, P., Goossens, M., Cornélis, F., Michou, L., Bardin, T. and Tchernitchko, D.O. 2007. The TNFRSF1A R92Q mutation is frequent in rheumatoid arthritis but shows no evidence for association or linkage with the disease. *Ann. Rheum. Dis.* 66: 1113-1115.
- Rossol, M., Meusch, U., Pierer, M., Kaltenhäuser, S., Häntzschel, H., Hauschildt, S. and Wagner, U. 2007. Interaction between transmembrane TNF and TNFR1/2 mediates the activation of monocytes by contact with T cells. *J. Immunol.* 179: 4239-4248.
- Kim, S., Moon, S.M., Kim, Y.S., Kim, J.J., Ryu, H.J., Kim, Y.J., Choi, J.W., Park, H.S., Kim, D.G., Shin, H.D., Rutherford, M.S., Oh, B. and Lee, J.K. 2008. TNFR1 promoter -329G/T polymorphism results in allele-specific repression of TNFR1 expression. *Biochem. Biophys. Res. Commun.* 368: 395-401.
- Paland, N., Böhme, L., Gurumurthy, R.K., Mäurer, A., Szczepek, A.J. and Rudel, T. 2008. Reduced display of tumor necrosis factor receptor I at the host cell surface supports infection with *Chlamydia trachomatis*. *J. Biol. Chem.* 283: 6438-6448.
- Cereda, C., Baiocchi, C., Bongioanni, P., Cova, E., Guareschi, S., Metelli, M.R., Rossi, B., Sbalsi, I., Cuccia, M.C. and Ceroni, M. 2008. TNF and sTNFR1/2 plasma levels in ALS patients. *J. Neuroimmunol.* 194: 123-131.
- Gupta, R., Sharma, S.C. and Das, S.N. 2008. Association of TNF- α and TNFR1 promoters and 3' UTR region of TNFR2 gene polymorphisms with genetic susceptibility to tobacco-related oral carcinoma in Asian Indians. *Oral Oncol.* 44: 455-463.

CHROMOSOMAL LOCATION

Genetic locus: Tnfrsf1a (mouse) mapping to 6 F3.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PRODUCT

TNF-R1 (m): 293T Lysate represents a lysate of mouse TNF-R1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

TNF-R1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TNF-R1 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

TNF-R1 (G-2): sc-374185 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse TNF-R1 expression in TNF-R1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



TNF-R1 (G-2): sc-374185. Western blot analysis of TNF-R1 expression in non-transfected: sc-117752 (A) and mouse TNF-R1 transfected: sc-124202 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.