



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TRMT1 (m): 293T Lysate: sc-124296

BACKGROUND

Transfer RNA (tRNA) modifications help regulate the efficiency of mRNA translation by maintaining the correct reading frames. N2,N2-dimethylguanosine tRNA methyltransferase, also known as TRMT1 or tRNA(guanine-26,N(2)-N(2)) methyltransferase, is a 659 amino acid enzyme that is responsible for tRNA modifications in eukaryotes. Using S-adenosyl-L-methionine as a methyl donor, TRMT1 dimethylates a single guanine residue at position 26 of tRNA. TRMT1, which was initially identified in yeast and *C. elegans*, has a 26% and 31% sequence identity to its yeast and *C. elegans* homologs, respectively. There are two isoforms of TRMT1 produced by alternative splicing events. The TRMT1 gene maps to chromosome 19p13.13 and mutations in this gene lead to abrogated enzyme activity and a decrease in protein levels.

REFERENCES

- Edqvist, J., Straby, K.B. and Grosjean, H. 1995. Enzymatic formation of N2,N2-dimethylguanosine in eukaryotic tRNA: importance of the tRNA architecture. *Biochimie* 77: 54-61.
- Constantinesco, F., Benachenhou, N., Motorin, Y. and Grosjean, H. 1998. The tRNA(guanine-26,N2-N2) methyltransferase (TRM1) from the hyperthermophilic archaeon *Pyrococcus furiosus*: cloning, sequencing of the gene and its expression in *Escherichia coli*. *Nucleic Acids Res.* 26: 3753-3761.
- Liu, J., Liu, J. and Straby, K.B. 1998. Point and deletion mutations eliminate one or both methyl group transfers catalysed by the yeast TRM1 encoded tRNA (m22G26)dimethyltransferase. *Nucleic Acids Res.* 26: 5102-5108.
- Björk, G.R., Durand, J.M., Hagervall, T.G., Leipuviene, R., Lundgren, H.K., Nilsson, K., Chen, P., Qian, Q. and Urbonavicius, J. 1999. Transfer RNA modification: influence on translational frameshifting and metabolism. *FEBS Lett.* 452: 47-51.
- Niederberger, C., Gräub, R., Costa, A., Desgrès, J. and Schweingruber, M.E. 1999. The tRNA N2,N2-dimethylguanosine-26 methyltransferase encoded by gene TRM1 increases efficiency of suppression of an ochre codon in *Schizosaccharomyces pombe*. *FEBS Lett.* 464: 67-70.
- Constantinesco, F., Motorin, Y. and Grosjean, H. 1999. Characterisation and enzymatic properties of tRNA(guanine 26, N2, N2)-dimethyltransferase (Trm1p) from *Pyrococcus furiosus*. *J. Mol. Biol.* 291: 375-392.
- Liu, J. and Straby, K.B. 2000. The human tRNA(m(2)(2)G(26))dimethyltransferase: functional expression and characterization of a cloned hTRM1 gene. *Nucleic Acids Res.* 28: 3445-3451.
- Urbonavicius, J., Qian, Q., Durand, J.M., Hagervall, T.G. and Björk, G.R. 2001. Improvement of reading frame maintenance is a common function for several tRNA modifications. *EMBO J.* 20: 4863-4873.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611669. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Trmt1 (mouse) mapping to 8 C3.

PRODUCT

TRMT1 (m): 293T Lysate represents a lysate of mouse TRMT1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

TRMT1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TRMT1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


TRMT1 (F-8): sc-514876 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse TRMT1 expression in TRMT1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG_x BP-HRP: sc-516102 or m-IgG_x BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



TRMT1 (F-8): sc-514876. Western blot analysis of TRMT1 expression in non-transfected: sc-117752 (**A**) and mouse TRMT1 transfected: sc-124296 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.