



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TrxR2 (m2): 293T Lysate: sc-124315

BACKGROUND

Thioredoxin (Trx) is a redox protein that is found in several species, such as bacteria, plants and mammals, and contains a conserved active site, consisting of Trp-Cys-Gly-Pro-Cys. Trx has several biological functions. It acts as a hydrogen donor for ribonucleotide reductase, which is critical for DNA synthesis, and modulates the DNA-binding activity of several transcription factors, including NFκB, AP-1, p53, TFIIIC and glucocorticoid receptor. Trx also stimulates cell growth, is an inhibitor of apoptosis and plays a role in the protection against oxidative stress. Drugs that inhibit Trx have antitumor activity, suggesting that Trx is involved in a variety of human diseases, including cancer. Thioredoxin 2 (Trx-2) is a small redox protein that is localized to the mitochondria and is essential for cell viability, playing a crucial role in the scavenging of ROS in mitochondria and regulating the mitochondrial apoptosis signaling pathway. Trx reductases (TrxR1 and TrxR2) are ubiquitously expressed flavoproteins that catalyze the NADPH-dependent reduction of Trx as well as several other oxidized cellular components. Mammalian Trx reductases are a part of a selenium-containing pyridine nucleotide-disulphide oxidoreductase family, which has a conserved catalytic site of Cys-Val-Asn-Val-Gly-Cys. TrxR1 and TrxR2 are also involved in the prevention of oxidative stress. Inhibition of TrxR activity may provide for potential treatments of cancer, AIDS and other autoimmune diseases, as well as bacterial infections and parasitic diseases.

REFERENCES

- Soderberg, A., et al. 1998. Monoclonal antibodies to human Thioredoxin reductase. *Biochem. Biophys. Res. Commun.* 249: 86-89.
- Lee, S.R., et al. 1999. Molecular cloning and characterization of a mitochondrial selenocysteine-containing Thioredoxin reductase from rat liver. *J. Biol. Chem.* 274: 4722-4734.
- Miranda-Vizuete, A., et al. 1999. Human mitochondrial Thioredoxin reductase cDNA cloning, expression and genomic organization. *Eur. J. Biochem.* 261: 405-412.
- Gorlatov, S.N. and Stadtman, T.C. 1999. Human selenium-dependent Thioredoxin reductase from HeLa cells: properties of forms with differing heparin affinities. *Arch. Biochem. Biophys.* 369: 133-142.
- Miranda-Vizuete, A., et al. 1999. cDNA cloning, expression and chromosomal localization of the mouse mitochondrial Thioredoxin reductase gene. *Biochim. Biophys. Acta* 1447: 113-118.
- Mustacich, D. and Powis, G. 2000. Thioredoxin reductase. *Biochem. J.* 346: 1-8.
- Conrad, M., et al. 2004. Essential role for mitochondrial Thioredoxin reductase in hematopoiesis, heart development, and heart function. *Mol. Cell. Biol.* 24: 9414-9423.
- Nalvarte, I., et al. 2004. Overexpression of enzymatically active human cytosolic and mitochondrial Thioredoxin reductase in HEK-293 cells. Effect on cell growth and differentiation. *J. Biol. Chem.* 279: 54510-54517.
- Kim, J.R., et al. 2004. Oxidation of Thioredoxin reductase in HeLa cells stimulated with tumor necrosis factor-α. *FEBS Lett.* 567: 189-196.

CHROMOSOMAL LOCATION

Genetic locus: Txnr2 (mouse) mapping to 16 A3.

PRODUCT

TrxR2 (m2): 293T Lysate represents a lysate of mouse TrxR2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

TrxR2 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive TrxR2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

TrxR2 (F-5): sc-376868 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse TrxR2 expression in TrxR2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



TrxR2 (F-5): sc-376868. Western blot analysis of TrxR2 expression in non-transfected: sc-117752 (A) and mouse TrxR2 transfected: sc-124315 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.