



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



UDP-GlcDH (m): 293T Lysate: sc-124442

BACKGROUND

UDP-GlcDH (also called UDP-glucose 6-dehydrogenase, UGDH or UDPGDH) is a member of the UDP-glucose/GDP-mannose dehydrogenase family. UDP-GlcDH converts UDP-glucose to UDP-glucuronic acid, which is a crucial component in the biosynthesis of the glycosaminoglycans, hyaluronan, heparan sulfate and chondroitin sulfate. Found as common components of the extracellular matrix, these glycosaminoglycans are significant in signal transduction, cell migration, cancer growth and cancer metastasis. UDP-glucuronic acid (UDP-GlcA) is needed in the liver for the excretion of toxic compounds. UDP-GlcDH is a ubiquitously expressed protein most abundant in the liver. The protein structure of UDP-GlcDH was first analyzed in cow liver and found to be a homohexamer. This structure is well conserved between species and phyla with an overall 97% sequence identity shared between different species of mammals. Research indicates that UDP-GlcDH expression is upregulated by TGF β and downregulated by hypoxia.

REFERENCES

1. Hempel, J., Perozich, J., Romovacek, H., Hinich, A., Kuo, I. and Feingold, D.S. 1994. UDP-glucose dehydrogenase from bovine liver: primary structure and relationship to other dehydrogenases. *Protein Sci.* 3: 1074-1080.
2. Spicer, A.P., Kaback, L.A., Smith, T.J. and Seldin, M.F. 1998. Molecular cloning and characterization of the human and mouse UDP-glucose dehydrogenase genes. *J. Biol. Chem.* 273: 25117-25124.
3. Marcu, O., Stathakis, D.G. and Marsh, J.L. 1999. Assignment of the UGDH locus encoding UDP-glucose dehydrogenase to human chromosome band 4p15.1 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 86: 244-245.
4. Johansson, H., Sterky, F., Amini, B., Lundeberg, J. and Kleczkowski, L.A. 2002. Molecular cloning and characterization of a cDNA encoding poplar UDP-glucose dehydrogenase, a key gene of hemicellulose/pectin formation. *Biochim. Biophys. Acta* 1576: 53-58.
5. Bontemps, Y., Vuillermoz, B., Antonicelli, F., Perreau, C., Danan, J.L., Maquart, F.X. and Wegrowski, Y. 2003. Specific protein-1 is a universal regulator of UDP-glucose dehydrogenase expression: its positive involvement in transforming growth factor- β signaling and inhibition in hypoxia. *J. Biol. Chem.* 278: 21566-21575.
6. Vatsyayan, J., Peng, H.L. and Chang, H.Y. 2005. Analysis of human UDP-glucose dehydrogenase gene promoter: identification of an Sp1 binding site crucial for the expression of the large transcript. *J. Biochem.* 137: 703-709.
7. Huh, J.W., Choi, M.M., Yang, S.J., Yoon, S.Y., Choi, S.Y. and Cho, S.W. 2005. Inhibition of human UDP-glucose dehydrogenase expression using siRNA expression vector in breast cancer cells. *Biotechnol. Lett.* 27: 1229-1232.
8. SWISS-PROT/TrEMBL (060701). World Wide Web URL:
<http://www.expasy.ch/sprot/sprot-top.html>
9. LocusLink Report (LocusID: 7358) <http://www.ncbi.nlm.nih.gov/LocusLink/>

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Ugdh (mouse) mapping to 5 C3.1.

PRODUCT

UDP-GlcDH (m): 293T Lysate represents a lysate of mouse UDP-GlcDH transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

UDP-GlcDH (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive UDP-GlcDH antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.