



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

UIP1 (m): 293T Lysate: sc-124465

BACKGROUND

UIP1 (UCHL5-interacting protein, HAUS augmin-like complex subunit 7) is a 368 amino acid product of the human HAUS7 gene. UIP1 is subunit 7 of the HAUS augmin-like complex, which is involved with regulation of spindle assembly during mitosis and contributes to the integrity of the centrosome. The HAUS augmin-like complex associates with the γ -tubulin ring complex. This interaction is required for spindle assembly. UIP1 also binds to UCHL5 (ubiquitin carboxyl-terminal hydrolase isozyme L5) which is a protease that specifically cleaves 'Lys-48'-linked polyubiquitin chains. UIP1 localizes to the centrosome during interphase and becomes part of the spindle microtubules during mitosis. Detectable levels of expression have been found in spleen, thymus, testis, ovary, small intestine and colon tissues.

REFERENCES

- Esposito, T., et al. 1997. Expressed STSs and transcription of human Xq28. *Gene* 187: 185-191.
- Li, T., et al. 2001. Identification of two proteins, S14 and UIP1, that interact with UCH37. *FEBS Lett.* 488: 201-205.
- Wang, B., et al. 2008. Evaluation of the low-specificity protease elastase for large-scale phosphoproteome analysis. *Anal. Chem.* 80: 9526-9533.
- Dephoure, N., et al. 2008. A quantitative atlas of mitotic phosphorylation. *Proc. Natl. Acad. Sci. USA* 105: 10762-10767.
- Lawo, S., et al. 2009. HAUS, the 8-subunit human Augmin complex, regulates centrosome and spindle integrity. *Curr. Biol.* 19: 816-826.
- Uehara, R., et al. 2009. The augmin complex plays a critical role in spindle microtubule generation for mitotic progression and cytokinesis in human cells. *Proc. Natl. Acad. Sci. USA* 106: 6998-7003.
- Mayya, V., et al. 2009. Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. *Sci. Signal.* 2: ra46.

CHROMOSOMAL LOCATION

Genetic locus: Haus7 (mouse) mapping to X A7.3.

PRODUCT

UIP1 (m): 293T Lysate represents a lysate of mouse UIP1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

UIP1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive UIP1 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

UIP1 (C-2): sc-393259 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse UIP1 expression in UIP1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

DATA



UIP1 (C-2): sc-393259. Western blot analysis of UIP1 expression in non-transfected: sc-117752 (A) and mouse UIP1 transfected: sc-124465 (B) 293T whole cell lysates.



UIP1 (E-13): sc-162379. Western blot analysis of UIP1 expression in non-transfected: sc-117752 (A) and mouse UIP1 transfected: sc-124465 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.