



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

WWOX (m): 293T Lysate: sc-124659

BACKGROUND

WWOX (WW domain containing oxidoreductase) protein is a candidate tumor suppressor consisting of two WW domains that influence protein-protein interactions, and a short chain dehydrogenase (SDR) domain, that influences sex-steroid metabolism. Modulation of the WWOX gene influences esophageal squamous cell carcinogenesis and tumorigenicity of breast cancer cell lines MDA-MB-435 and T47D. The murine homolog WOX1 localizes in the mitochondria, and contains a mitochondrial targeting sequence mapping within the SDR domain. JNK1 can physically associate with WOX1 and sequester WOX1-dependent apoptosis.

REFERENCES

1. Bednarek, A.K., et al. 2001. WWOX, the FRA16D gene, behaves as a suppressor of tumor growth. *Cancer Res.* 61: 8068-8073.
2. Chang, N.S., et al. 2001. Hyaluronidase induction of a WW domain-containing oxidoreductase that enhances tumor necrosis factor cytotoxicity. *J. Biol. Chem.* 276: 3361-3370.
3. Kuroki, T., et al. 2002. Genetic alterations of the tumor suppressor gene WWOX in esophageal squamous cell carcinoma. *Cancer Res.* 62: 2258-2260.
4. Ludes-Meyers, J.H., et al. 2003. WWOX, the common chromosomal fragile site, FRA16D, cancer gene. *Cytogenet Genome Res.* 100: 101-110.
5. Chang, N.S., et al. 2003. JNK1 physically interacts with WW domain-containing oxidoreductase (WOX1) and inhibits WOX1-mediated apoptosis. *J. Biol. Chem.* 278: 9195-9202.

CHROMOSOMAL LOCATION

Genetic locus: *Wwox* (mouse) mapping to 8 E1.

PRODUCT

WWOX (m): 293T Lysate represents a lysate of mouse WWOX transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

WWOX (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive WWOX antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

WWOX (C-7): sc-374449 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse WWOX expression in WWOX transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



WWOX (C-7): sc-374449. Western blot analysis of WWOX expression in non-transfected: sc-117752 (A) and mouse WWOX transfected: sc-124659 (B) 293T whole cell lysates.



WWOX (A-5): sc-373846. Western blot analysis of WWOX expression in non-transfected: sc-117752 (A) and mouse WWOX transfected: sc-124659 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.