



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ZNF622 (m): 293T Lysate: sc-124806

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZNF622 (zinc-finger protein 622), also known as ZPR9 (zinc-finger-like protein 9), is a 477 amino acid protein that localizes to both the nucleus and the cytoplasm. Expressed in liver, spleen, lung, kidney and brain, ZNF622 is thought to activate the bound transcription factor B-Myb and, through this activation, may play a role in embryonic development. ZNF622 contains two U1-type zinc fingers and exists as either a homodimer or a heterodimer that can be phosphorylated by MELK (maternal embryonic leucine zipper kinase). Overexpression of ZNF622 may be associated with liver metastases, carcinomas and colorectal carcinomas.

## REFERENCES

- Thiesen, H.J. 1990. Multiple genes encoding zinc-finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
- Huebner, K., Druck, T., Croce, C.M. and Thiesen, H.J. 1991. Twenty-seven non-overlapping zinc-finger cDNAs from human T cells map to nine different chromosomes with apparent clustering. *Am. J. Hum. Genet.* 48: 726-740.
- Seong, H.A., Gil, M., Kim, K.T., Kim, S.J. and Ha, H. 2002. Phosphorylation of a novel zinc-finger-like protein, ZPR9, by murine protein serine/threonine kinase 38 (MPK38). *Biochem. J.* 361 (Pt 3): 597-604.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608694. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Seong, H.A., Kim, K.T. and Ha, H. 2003. Enhancement of B-Myb transcriptional activity by ZPR9, a novel zinc-finger protein. *J. Biol. Chem.* 278: 9655-9662.
- Andersen, J.S., Lam, Y.W., Leung, A.K., Ong, S.E., Lyon, C.E., Lamond, A.I. and Mann, M. 2005. Nucleolar proteome dynamics. *Nature* 433: 77-83.
- Kleivi, K., Lind, G.E., Diep, C.B., Meling, G.I., Brandal, L.T., Nesland, J.M., Myklebost, O., Rognum, T.O., Giercksky, K.E., Skotheim, R.I. and Lothe, R.A. 2007. Gene expression profiles of primary colorectal carcinomas, liver metastases, and carcinomas. *Mol. Cancer* 6: 2.

## CHROMOSOMAL LOCATION

Genetic locus: Zfp622 (mouse) mapping to 15 B1.

## PRODUCT

ZNF622 (m): 293T Lysate represents a lysate of mouse ZNF622 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

ZNF622 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ZNF622 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ZNF622 (3I4): sc-100980 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse ZNF622 expression in ZNF622 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



ZNF622 (3I4): sc-100980. Western blot analysis of ZNF622 expression in non-transfected: sc-117752 (A) and mouse ZNF622 transfected: sc-124806 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.