



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



α Enolase (m6): 293T Lysate: sc-124902

BACKGROUND

Enolases have been characterized as highly conserved cytoplasmic glycolytic enzymes that may be involved in differentiation. Three isoenzymes have been identified, α Enolase, β Enolase and γ Enolase. α Enolase expression has been detected on most tissues, whereas β Enolase is expressed predominantly in muscle tissue and γ Enolase is detected only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphoenolpyruvic acid in the glycolytic pathway.

REFERENCES

- Whitehead, M.C., Marangos, P.J., Connolly, S.M. and Morest, D.K. 1982. Synapse formation is related to the onset of neuron-specific Enolase immunoreactivity in the avian auditory and vestibular systems. *Dev. Neurosci.* 5: 298-307.
- Verma, M. and Dutta, S.K. 1994. DNA sequences encoding Enolase are remarkably conserved from yeast to mammals. *Life Sci.* 55: 893-899.
- Keller, A., Bérod, A., Dussaillant, M., Lemandé, N., Gros, F. and Lucas, M. 1994. Coexpression of α and γ Enolase genes in neurons of adult rat brain. *J. Neurosci. Res.* 38: 493-504.
- Zhang, E., Brewer, J.M., Minor, W., Carreira, L.A. and Lebioda, L. 1997. Mechanism of Enolase: the crystal structure of asymmetric dimer Enolase-2-phospho-D glycerate/Enolase-phosphoenolpyruvate at 2.0 Å resolution. *Biochemistry* 36: 12526-12534.
- Deloulme, J.C., Helies, A., Ledig, M., Lucas, M. and Sensenbrenner, M. 1997. A comparative study of the distribution of α and γ Enolase subunits in cultured rat neural cells and fibroblasts. *Int. J. Dev. Neurosci.* 15: 183-194.
- Sensenbrenner, M., Lucas, M. and Deloulme, J.C. 1997. Expression of two neuronal markers, growth-associated protein 43 and neuron-specific Enolase, in rat glial cells. *J. Mol. Med.* 75: 653-663.
- Ito, S., Honma, T., Ishida, K., Wada, N., Sasaoka, S., Hosoda, M. and Nohno, T. 2007. Differential expression of the human α Enolase gene in oral epithelium and squamous cell carcinoma. *Cancer Sci.* 98: 499-505.
- Perconti, G., Ferro, A., Amato, F., Rubino, P., Randazzo, D., Wolff, T., Feo, S. and Giallongo, A. 2007. The kelch protein NS1-BP interacts with α Enolase/Mbp1 and is involved in c-Myc gene transcriptional control. *Biophys. Acta* 1773: 1774-1785.

CHROMOSOMAL LOCATION

Genetic locus: Eno1 (mouse) mapping to 4 E2.

PRODUCT

α Enolase (m6): 293T Lysate represents a lysate of mouse α Enolase transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

α Enolase (m6): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive α Enolase antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.