



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AKR7A5 (m): 293T Lysate: sc-124941

BACKGROUND

Aldo-keto reductase 7 (AKR7) functions in the metabolism of aflatoxin B1 and other dicarbonyl-containing compounds with ketone groups on adjacent carbon atoms in a broad range of tissues, notably the liver. The AKR7 gene maps to human chromosome 1p36.13, a region frequently deleted in sporadic colorectal cancer. The functional significance of this correlation lies in the constitutive expression of AKR7 in human liver to eliminate aflatoxin (an environmental carcinogen), thus acting as an endogenous chemo-preventative agent.

REFERENCES

1. Ellis, E.M., et al. 1995. Substrate specificity of an aflatoxin-metabolizing aldehyde reductase. *Biochem. J.* 312: 535-541.
2. Ireland, L.S., et al. 1998. Molecular cloning, expression and catalytic activity of a human AKR7 member of the aldo-keto reductase superfamily: evidence that the major 2-carboxybenzaldehyde reductase from human liver is a homologue of rat aflatoxin B1-aldehyde reductase. *Biochem. J.* 332: 21-34.
3. Kelly, V.P., et al. 2000. Purification from rat liver of a novel constitutively expressed member of the aldo-keto reductase 7 family that is widely distributed in extrahepatic tissues. *Biochem. J.* 348: 389-400.
4. Kelly, V.P., et al. 2002. Novel homodimeric and heterodimeric rat γ -hydroxybutyrate synthases that associate with the Golgi apparatus define a distinct subclass of aldo-keto reductase 7 family proteins. *Biochem. J.* 366: 847-861.
5. Praml, C., et al. 2003. Aflatoxin B1 aldehyde reductase (AFAR) genes cluster at 1p35-1p36.1 in a region frequently altered in human tumour cells. *Oncogene* 22: 4765-4773.
6. Hyndman, D., et al. 2003. The aldo-keto reductase superfamily homepage. *Chem Biol Interact.* 143-144: 621-631.
7. Kozma, E., et al. 2003. The high resolution crystal structure of rat liver AKR7A1: understanding the substrate specificities of the AKR7 family. *Chem Biol Interact.* 143-144: 289-297.
8. Grant, A.W., et al. 2003. A novel aldo-keto reductase from *Escherichia coli* can increase resistance to methylglyoxal toxicity. *FEMS Microbiol Lett.* 218: 93-99.
9. Gardner, R., et al. 2004. Detoxication of the environmental pollutant acrolein by a rat liver aldo-keto reductase. *Toxicol Lett.* 148: 65-72.

CHROMOSOMAL LOCATION

Genetic locus: *Akr7a5* (mouse) mapping to 4 D3.

PRODUCT

AKR7A5 (m): 293T Lysate represents a lysate of mouse AKR7A5 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

AKR7A5 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive AKR7A5 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.