



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CD72 (m): 293T Lysate: sc-125116

BACKGROUND

CD5 has been identified as a transmembrane glycoprotein that is expressed on 70% of normal peripheral blood lymphocytes and on virtually all T lymphocytes in thymus and peripheral blood. Activation of T cells through the T cell receptor (TCR) results in tyrosine phosphorylation of CD5, and the absence of CD5 renders T cells hyper-responsive to TCR-mediated activation. CD5 associates with the TCR/CD3 ζ chain, and with the Src family kinase, Lck p56. The C-type lectin superfamily member CD72 is a cell surface negative regulator of B cell activation from the pro-B through the mature B cell stage. CD72 serves as a receptor for CD5. The ability of lymphocytes to respond to antigenic or mitogenic stimulation utilizes both positive and negative regulatory proteins that influence the threshold for responsiveness. The human CD72 gene maps to chromosome 9p13.3 and encodes a transmembrane glycoprotein that contains an immunoreceptor tyrosine-based inhibition motif (ITIM). Upon tyrosine phosphorylation, the CD72 ITIM recruits SH2-containing phosphatases such as SHP-1, resulting in downregulation of cell activation. CD72^{-/-} mice contain hyperproliferative B cells.

REFERENCES

1. Knapp, W., et al, eds. 1989. Leucocyte Typing IV. New York: Oxford University Press.
2. Schlossman, S., et al, eds. 1995. Leucocyte Typing V. Oxford: Oxford University Press.
3. Ying, H., et al. 1998. Regulation of mouse CD72 gene expression during B lymphocyte development. *J. Immunol.* 161: 4760-4767.
4. Wu, Y., et al. 1998. The B cell transmembrane protein CD72 binds to and is an *in vivo* substrate of the protein tyrosine phosphatase SHP-1. *Curr. Biol.* 8: 1009-1017.
5. Pan, C., et al. 1999. CD72-deficient mice reveal nonredundant roles of CD72 in B cell development and activation. *Immunity* 11: 495-506.
6. Parnes, J.R., et al. 2000. CD72, a negative regulator of B cell responsiveness. *Immunol. Rev.* 176: 75-85.
7. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 107272. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Wu, H.J., et al. 2002. Positive and negative roles of CD72 in B cell function. *Immunol. Res.* 25: 155-166.

CHROMOSOMAL LOCATION

Genetic locus: Cd72 (mouse) mapping to 4 B1.

PRODUCT

CD72 (m): 293T Lysate represents a lysate of mouse CD72 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

CD72 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CD72 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.