



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# CD79B (m): 293T Lysate: sc-125117

## BACKGROUND

CD79 (also designated Ig  $\alpha$ /Ig  $\beta$ ) is a heterodimer composed of  $\alpha$  chains, designated CD79A or MB-1, and  $\beta$  chains, designated CD79B or B29. The B cell antigen receptor complex (BCR) is formed by the association of CD79 with a membrane immunoglobulin, such as IgM or IgD. The membrane immunoglobulins IgM and IgD achieve surface expression and antigen presentation function in response to CD79 association. The cytoplasmic tails of both CD79A and CD79B contain an ITAM (immunoreceptor tyrosine-based activation) motif, which acts to initiate the BCR signaling reactions by binding to and activating tyrosine kinases.

## REFERENCES

1. Poppema, S., et al. 1987. Monoclonal antibodies (MT1, MT2, MB1, MB2, MB3) reactive with leukocyte subsets in paraffin-embedded tissue sections. Am. J. Pathol. 127: 418-429.
2. van Noesel, C.J., et al. 1991. The membrane IgM-associated heterodimer on human B cells is a newly defined B cell antigen that contains the protein product of the mb-1 gene. J. Immunol. 146: 3881-3888.
3. Mason, D.Y., et al. 1991. The IgM-associated protein mb-1 as a marker of normal and neoplastic B cells. J. Immunol. 147: 2474-2482.
4. Ha, H.J., et al. 1992. Molecular cloning and expression pattern of a human gene homologous to the murine mb-1 gene. J. Immunol. 148: 1526-1531.
5. Mason, D.Y., et al. 1992. The B29 and mb-1 polypeptides are differentially expressed during human B cell differentiation. Eur. J. Immunol. 22: 2753-2756.
6. Jones, M., et al. 1993. Detection of T and B cells in many animal species using cross-reactive anti-peptide antibodies. J. Immunol. 150: 5429-5435.
7. Wood, W.J., Jr., et al. 1993. Isolation and chromosomal mapping of the human immunoglobulin-associated B29 gene (IGB). Genomics 16: 187-192.
8. Mason, D.Y., et al. 1995. CD79A: a novel marker for B-cell neoplasms in routinely processed tissue samples. Blood 86: 1453-1459.
9. Macardle, P.J., et al. 1997. The antigen receptor complex on cord B lymphocytes. Immunology 90: 376-382.

## CHROMOSOMAL LOCATION

Genetic locus: Cd79b (mouse) mapping to 11 E1.

## PRODUCT

CD79B (m): 293T Lysate represents a lysate of mouse CD79B transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

CD79B (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CD79B antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


CD79B (B29/123): sc-53210 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse CD79B expression in CD79B transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG $\lambda$  BP-HRP: sc-516132 or m-IgG $\lambda$  BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



CD79B (B29/123): sc-53210. Western blot analysis of CD79B expression in non-transfected: sc-117752 (**A**) and mouse CD79B transfected: sc-125117 (**B**) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.