



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CTRP1 (m): 293T Lysate: sc-125181

BACKGROUND

CTRP1 (also designated complement C1q tumor necrosis factor-related protein 1 precursor, G protein-coupled receptor-interacting protein and GIP) is a 281 amino acid protein encoded by the human gene C1QTN1. CTRP1 is a member of the CTRP superfamily and is expressed at high levels in adipose tissues of obese Zucker diabetic fatty (fa/fa) rats. It consists of an N-terminal signal peptide sequence followed by a variable region, a collagen repeat domain and a C-terminal globular domain. CTRP1 expression is induced by proinflammatory cytokines, including TNF α and IL-1 β . Due to its cytokine induced nature, CTRP1 expression may be associated with a low-grade chronic inflammation status in adipose tissues.

REFERENCES

1. Wu, T.H., Wu, S.C., Huang, T.P., Yu, C.L. and Tsai, C.Y. 1997. Increased excretion of tumor necrosis factor α and interleukin 1 β in urine from patients with IgA nephropathy and Schönlein-Henoch purpura. *Nephron* 74: 79-88.
2. Branten, A.J., Klasen, I.S. and Wetzels, J.F. 2002. Short-term effects of fish oil treatment on urinary excretion of high- and low-molecular weight proteins in patients with IgA nephropathy. *Clin. Nephrol.* 58: 267-274.
3. Bergmann, J., Buchheidt, D., Waldherr, R., Maywald, O., van der Woude, F.J., Hehlmann, R. and Braun, C. 2005. IgA nephropathy and hodgkin's disease: a rare coincidence. Case report and literature review. *Am. J. Kidney Dis.* 45: e16-e19.
4. Kim, K.Y., Kim, H.Y., Kim, J.H., Lee, C.H., Kim, D.H., Lee, Y.H., Han, S.H., Lim, J.S., Cho, D.H., Lee, M.S., Yoon, S., Kim, K.I., Yoon, D.Y. and Yang, Y. 2006. Tumor necrosis factor- α and interleukin-1 β increases CTRP1 expression in adipose tissue. *FEBS Lett.* 580: 3953-3960.
5. Khositseth, S., Kanitsap, N., Warnnissorn, N. and Thongboonkerd, V. 2007. IgA nephropathy associated with Hodgkin's disease in children: a case report, literature review and urinary proteome analysis. *Pediatr. Nephrol.* 22: 541-546.
6. Jeon, J.H., Kim, K.Y., Kim, J.H., Baek, A., Cho, H., Lee, Y.H., Kim, J.W., Kim, D., Han, S.H., Lim, J.S., Kim, K.I., Yoon, D.Y., Kim, S.H., Oh, G.T., Kim, E. and Yang, Y. 2008. A novel adipokine CTRP1 stimulates aldosterone production. *FASEB J.* 22: 1502-1511.

CHROMOSOMAL LOCATION

Genetic locus: C1qtnf1 (mouse) mapping to 11 E2.

PRODUCT

CTRP1 (m): 293T Lysate represents a lysate of mouse CTRP1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20 $^{\circ}$ C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

CTRP1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CTRP1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.