



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



FUSIP1 (m2): 293T Lysate: sc-125352



The Power to Question

BACKGROUND

FUSIP1 (FUS interacting protein (serine/arginine-rich) 1), also known as NSSR, TASR (TLS-associated protein with Ser-Arg repeats), SRP38, TASR1, TASR2, FUSIP2, SFRS13 or SRP40 (40 kDa SR-repressor protein), is a member of the serine/arginine (SR) family of splicing factors. Members of the SR family all contain one or more RNA recognition motifs (RRM) and an SR-rich domain. SR factors are not only essential for constitutive splicing but also regulate splicing in a concentration-dependent manner by influencing the selection of alternative splice sites. Expressed in a variety of tissues with low expression in kidney, liver and heart, FUSIP1 localizes to the cytoplasm and nuclear speckles. In its dephosphorylated form (occurring during M phase of the cell cycle), FUSIP1 functions as a potent general repressor of pre-mRNA splicing and can interact with U1 SnRNP 70. In its phosphorylated form, FUSIP1 interacts with Tra-2 β and, together, they may cooperate in the regulation of splicing. Four isoforms exist for FUSIP1. In neurons, FUSIP1 isoforms may act to either positively or negatively regulate alternative splicing.

REFERENCES

1. Yang, L., Embree, L.J., Tsai, S. and Hickstein, D.D. 1998. Oncoprotein TLS interacts with serine-arginine proteins involved in RNA splicing. *J. Biol. Chem.* 273: 27761-27764.
2. Komatsu, M., Kominami, E., Arahata, K. and Tsukahara, T. 1999. Cloning and characterization of two neural-salient serine/arginine-rich (NSSR) proteins involved in the regulation of alternative splicing in neurones. *Genes Cells* 4: 593-606.
3. Shin, C. and Manley, J.L. 2002. The SR protein SRP38 represses splicing in M phase cells. *Cell* 111: 407-417.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605221. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Liu, L., Chen, X.H., Huang, J., Lin, J.J., Lin, W.M. and Xu, P. 2004. NSSR1 promotes neuronal differentiation of mouse embryonic carcinoma P19 cells. *Neuroreport* 15: 823-828.
6. Fushimi, K., Osumi, N. and Tsukahara, T. 2005. NSSRs/TASRs/SRP38s function as splicing modulators via binding to pre-mRNAs. *Genes Cells* 10: 531-541.
7. Matsushita, H., Blackburn, M.L., Klineberg, E., Zielinska-Kwiatkowska, A., Bolander, M.E., Sarkar, G., Suva, L.J., Chansky, H.A. and Yang, L. 2007. TASR-1 regulates alternative splicing of collagen genes in chondrogenic cells. *Biochem. Biophys. Res. Commun.* 356: 411-417.
8. Shi, Y. and Manley, J.L. 2007. A complex signaling pathway regulates SRP38 phosphorylation and pre-mRNA splicing in response to heat shock. *Mol. Cell* 28: 79-90.
9. Xiao, P.J., Hu, L., Li, J., Lin, W., Chen, X. and Xu, P. 2007. NSSR1 is regulated in testes development and cryptorchidism and promotes the exon 5-included splicing of CREB transcripts. *Mol. Reprod. Dev.* 74: 1363-1372.

CHROMOSOMAL LOCATION

Genetic locus: Fusip1 (mouse) mapping to 4 D3.

PRODUCT

FUSIP1 (m2): 293T Lysate represents a lysate of mouse FUSIP1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

FUSIP1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive FUSIP1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.