



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Med8 (m): 293T Lysate: sc-125596

## BACKGROUND

In mammalian cells, transcription is regulated in part by high molecular weight co-activating complexes that mediate signals between transcriptional activators and RNA polymerase II (Pol II). The mediator complex is one such multi-protein structure that functions as a bridge between regulatory proteins and Pol II, thereby regulating Pol II-dependent transcription. Med8 (mediator complex subunit 8), also known as arc32 (activator-recruited cofactor 32 kDa component), is a 268 amino acid protein that localizes to the nucleus and exists as a component of the mediator complex. Involved in the pathway of protein modification and ubiquitination, Med8 is involved in transcriptional regulation and may also recruit E3 ubiquitin-protein ligase complexes to proteins targeted for proteasomal degradation. Multiple isoforms of Med8 exist due to alternative splicing events.

## REFERENCES

- Jiang, Y.W., et al. 1998. Mammalian mediator of transcriptional regulation and its possible role as an end-point of signal transduction pathways. *Proc. Natl. Acad. Sci. USA* 95: 8538-8543.
- Näär, A.M., et al. 1999. Composite co-activator ARC mediates chromatin-directed transcriptional activation. *Nature* 398: 828-832.
- Brower, C.S., et al. 2002. Mammalian mediator subunit mMed8 is an Elongin BC-interacting protein that can assemble with CUL-2 and Rbx1 to reconstitute a ubiquitin ligase. *Proc. Natl. Acad. Sci. USA* 99: 10353-10358.
- Sato, S., et al. 2003. Identification of mammalian mediator subunits with similarities to yeast Mediator subunits Srb5, Srb6, Med11, and Rox3. *J. Biol. Chem.* 278: 15123-15127.
- Sato, S., et al. 2003. A mammalian homolog of *Drosophila melanogaster* transcriptional coactivator intersex is a subunit of the mammalian mediator complex. *J. Biol. Chem.* 278: 49671-49674.
- Tomomori-Sato, C., et al. 2004. A mammalian mediator subunit that shares properties with *Saccharomyces cerevisiae* mediator subunit Cse2. *J. Biol. Chem.* 279: 5846-5851.
- Sato, S., et al. 2004. A set of consensus mammalian mediator subunits identified by multidimensional protein identification technology. *Mol. Cell* 14: 685-691.
- Léonard, V.H., et al. 2006. Interaction of Bunyamwera Orthobunyavirus NSs protein with mediator protein MED8: a mechanism for inhibiting the interferon response. *J. Virol.* 80: 9667-9675.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 607956. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Med8 (mouse) mapping to 4 D2.1.

## PRODUCT

Med8 (m): 293T Lysate represents a lysate of mouse Med8 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

Med8 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Med8 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Med8 (A-3): sc-365713 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Med8 expression in Med8 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Med8 (A-3): sc-365713. Western blot analysis of Med8 expression in non-transfected: sc-117752 (A) and mouse Med8 transfected: sc-125596 (B) 293T whole cell lysates.



Med8 (A-5): sc-365960. Western blot analysis of Med8 expression in non-transfected: sc-117752 (A) and mouse Med8 transfected: sc-125596 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.