



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# MUP1 (m2): 293T Lysate: sc-125660



The Power to Question

## BACKGROUND

The major urinary proteins (MUPs) represent a family of androgen-regulated murine proteins that are synthesized in the liver, secreted into the bloodstream and excreted into the urine. MUP1 (major urinary protein 1), also known as Mup7, Up-1 or Ltn-1, is a 180 amino acid secreted murine protein that belongs to the calycin superfamily. Excreted at high levels in the urine of adult male mice, MUP1 functions to bind pheromones that are released from drying urine, thereby activating the pheromones and affecting the sexual behavior of females. Additionally, MUP1, along with other MUP proteins, may mediate male-male aggression through the accessory olfactory neural pathway.

## REFERENCES

1. Berger, F.G. and Szoka, P. 1981. Biosynthesis of the major urinary proteins in mouse liver: a biochemical genetic study. *Biochem. Genet.* 19: 1261-1273.
2. Krauter, K., Leinwand, L., D'Eustachio, P., Ruddle, F. and Darnell, J.E. 1982. Structural genes of the mouse major urinary protein are on chromosome 4. *J. Cell Biol.* 94: 414-417.
3. Held, W.A., Gallagher, J.F., Hohman, C.M., Kuhn, N.J., Sampsel, B.M. and Hughes, R.G. 1987. Identification and characterization of functional genes encoding the mouse major urinary proteins. *Mol. Cell. Biol.* 7: 3705-3712.
4. Shahani, K., Gilmartin, M. and Derman, E. 1987. Nucleotide sequences of liver, lachrymal, and submaxillary gland mouse major urinary protein mRNAs: mosaic structure and construction of panels of gene-specific synthetic oligo-nucleotide probes. *Mol. Cell. Biol.* 7: 1938-1946.
5. Lücke, C., Franzoni, L., Abbate, F., Löhr, F., Ferrari, E., Sorbi, R.T., Rüterjans, H. and Spisni, A. 1999. Solution structure of a recombinant mouse major urinary protein. *Eur. J. Biochem.* 266: 1210-1218.
6. Abbate, F., Franzoni, L., Löhr, F., Lücke, C., Ferrari, E., Sorbi, R.T., Rüterjans, H. and Spisni, A. 1999. Complete 1H, 15N and 13C assignment of a recombinant mouse major urinary protein. *J. Biomol. NMR* 15: 187-188.
7. Valkova, N., Yunis, R., Mak, S.K., Kang, K. and Kültz, D. 2005. Nek8 mutation causes overexpression of galectin-1, sorcin, and vimentin and accumulation of the major urinary protein in renal cysts of jck mice. *Mol. Cell. Proteomics* 4: 1009-1018.
8. Chamero, P., Marton, T.F., Logan, D.W., Flanagan, K., Cruz, J.R., Saghatelian, A., Cravatt, B.F. and Stowers, L. 2007. Identification of protein pheromones that promote aggressive behaviour. *Nature* 450: 899-902.
9. Logan, D.W., Marton, T.F. and Stowers, L. 2008. Species specificity in major urinary proteins by parallel evolution. *PLoS ONE* 3: e3280.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Mup1 (mouse) mapping to 4 B3.

## PRODUCT

MUP1 (m2): 293T Lysate represents a lysate of mouse MUP1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

MUP1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MUP1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.