



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



NAT-9 (m): 293T Lysate: sc-125690

BACKGROUND

Acetyltransferases and deacetylases are protein groups most often associated with oncogenesis and cell cycle regulation. NAT-9 (N-acetyltransferase 9), also known as EBSP (embryo brain-specific protein), is a 207 amino acid protein belonging to the acetyltransferase family and the GNAT subfamily. Containing a N-acetyltransferase domain, NAT-9 may be associated with psoriasis and psoriatic arthritis, a type of inflammatory/autoimmune disease that affects skin, tendons and/or joints of the hands and feet. Expressed as two isoforms produced by alternative splicing events, NAT-9 is encoded by a gene located on human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

REFERENCES

1. Helms, C., Cao, L., Krueger, J.G., Wijsman, E.M., Chamian, F., Gordon, D., Heffernan, M., Daw, J.A., Robarge, J., Ott, J., Kwok, P.Y., Menter, A. and Bowcock, A.M. 2003. A putative RUNX1 binding site variant between SLC9A3R1 and NAT-9 is associated with susceptibility to psoriasis. *Nat. Genet.* 35: 349-356.
2. Bowcock, A.M. and Cookson, W.O. 2004. The genetics of psoriasis, psoriatic arthritis and atopic dermatitis. *Hum. Mol. Genet.* 13: R43-R55.
3. Bowcock, A.M. 2005. The genetics of psoriasis and autoimmunity. *Annu. Rev. Genomics Hum. Genet.* 6: 93-122.
4. Yamada, R. and Ymamoto, K. 2005. Recent findings on genes associated with inflammatory disease. *Mutat. Res.* 573: 136-151.
5. Morar, N., Bowcock, A.M., Harper, J.I., Cookson, W.O. and Moffatt, M.F. 2006. Investigation of the chromosome 17q25 PSORS2 locus in atopic dermatitis. *J. Invest. Dermatol.* 126: 603-606.
6. Filer, C.E., Ho, P., Bruce, I.N., Worthington, J. and Barton, A. 2009. Investigation of association of genes NAT-9, SLC9A3R1 and Raptor on chromosome 17q25 with psoriatic arthritis. *Ann. Rheum. Dis.* 68: 292-293.
7. Danik, J.S., Paré, G., Chasman, D.I., Zee, R.Y., Kwiatkowski, D.J., Parker, A., Miletich, J.P. and Ridker, P.M. 2009. Novel loci, including those related to Crohn Disease, psoriasis, and inflammation, identified in a genome-wide association study of Fibrinogen in 17,686 women: the women's genome health study. *Circ. Cardiovasc. Genet.* 2: 134-141.

CHROMOSOMAL LOCATION

Genetic locus: NAT9 (human) mapping to 17q25.1.

PRODUCT

NAT-9 (m): 293T Lysate represents a lysate of mouse NAT-9 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

NAT-9 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive NAT-9 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.