



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PP2A-C α (m): 293T Lysate: sc-125847

BACKGROUND

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit and a catalytic subunit. Four major families of protein phosphatase catalytic subunits have been identified, designated PP1, PP2A, PP2B (calcineurin) and PP2C. Characteristic of the protein phosphatase complexes, the PP2A phosphatase core enzyme is composed of a regulatory subunit and a catalytic subunit, the latter of which exists as two isoforms, designated PP2A α and PP2A β . The multiple subunits of PP2A work in concert to regulate a variety of metabolic pathways, including transcription, translation, cell cycle progression and oncogenic transformation.

REFERENCES

1. Strack, S., et al. 2002. Protein phosphatase 2A holoenzyme assembly: identification of contacts between B-family regulatory and scaffolding A subunits. *J. Biol. Chem.* 277: 20750-20755.
2. Avdi, N.J., et al. 2002. A role for protein phosphatase-2A in p38 mitogen-activated protein kinase-mediated regulation of the c-Jun NH₂-terminal kinase pathway in human neutrophils. *J. Biol. Chem.* 277: 40687-40696.
3. Zhou, J., et al. 2003. Characterization of the A α and A β subunit isoforms of protein phosphatase 2A: differences in expression, subunit interaction, and evolution. *Biochem. J.* 369: 387-398.
4. Shern, J.F., et al. 2003. Cytosolic ARL2 is complexed with cofactor D and protein phosphatase 2A. *J. Biol. Chem.* 278: 40829-40836.
5. Prickett, T.D. and Brautigam, D.L. 2004. Overlapping binding sites in protein phosphatase 2A for association with regulatory A and α 4 (mTAP42) subunits. *J. Biol. Chem.* 279: 38912-38920.
6. Gergs, U., et al. 2004. Overexpression of the catalytic subunit of protein phosphatase 2A impairs cardiac function. *J. Biol. Chem.* 279: 40827-40834.
7. Avci, C.B., et al. 2007. Protein phosphatase 2A (PP2A) has a potential role in CAPE-induced apoptosis of CCRF-CEM cells via effecting human telomerase reverse transcriptase activity. *Hematology* 12: 519-525.
8. Sun, L., et al. 2008. EB1 promotes Aurora B kinase activity through blocking its inactivation by protein phosphatase 2A. *Proc. Natl. Acad. Sci. USA* 105: 7153-7158.

CHROMOSOMAL LOCATION

Genetic locus: Ppp2ca (mouse) mapping to 11 B1.3.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

PP2A-C α (m): 293T Lysate represents a lysate of mouse PP2A-C α transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

PP2A-C α (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PP2A-C α antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PP2A-C α / β (1D6): sc-80665 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse PP2A-C α expression in PP2A-C α transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.