



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Stat3 (m): 293T Lysate: sc-126053

## BACKGROUND

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of JAK kinases, which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- $\alpha$  and form a heterodimer, which is part of the ISGF-3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 $\beta$  appears to be activated by both while Stat3 $\alpha$  is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by Prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

## REFERENCES

- Zhong, Z., et al. 1994. Stat3: a Stat family member activated by tyrosine phosphorylation in response to epidermal growth factor and interleukin-6. *Science* 264: 95-98.
- Darnell, J.E., et al. 1994. JAK-Stat pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* 264: 1415-1421.
- Hou, J., et al. 1994. An interleukin-4-induced transcription factor: IL-4 Stat. *Science* 265: 1701-1706.
- Yamamoto, K., et al. 1994. Stat4, a novel  $\gamma$  interferon activation site-binding protein expressed in early myeloid differentiation. *Mol. Cell. Biol.* 14: 4342-4349.
- Pallard, C., et al. 1995. Interleukin-3, erythropoietin, and Prolactin activate a Stat5-like factor in lymphoid cells. *J. Biol. Chem.* 270: 15942-15945.
- Qureshi, S.A., et al. 1995. Tyrosine-phosphorylated Stat1 and Stat2 plus a 48 kDa protein all contact DNA in forming interferon-stimulated-gene factor 3. *Proc. Natl. Acad. Sci. USA* 92: 3829-3833.
- Schindler, C., et al. 1995. transcriptional responses to polypeptide ligands: the JAK-Stat pathway. *Annu. Rev. Biochem.* 64: 621-651.

## CHROMOSOMAL LOCATION

Genetic locus: Stat3 (mouse) mapping to 11 D.

## PRODUCT

Stat3 (m): 293T Lysate represents a lysate of mouse Stat3 transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## APPLICATIONS

Stat3 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Stat3 antibodies. Recommended use: 10-20  $\mu$ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Stat3 (F-2): sc-8019 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Stat3 expression in Stat3 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Stat3 (F-2): sc-8019. Western blot analysis of Stat3 expression in non-transfected: sc-117752 (A) and mouse Stat3 transfected: sc-126053 (B) 293T whole cell lysates.



p-Stat3 (9E12): sc-81523. Western blot analysis of Stat3 phosphorylation in non-transfected: sc-117752 (A) and mouse Stat3 transfected: sc-126053 (B) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.