



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# STC2 (m): 293T Lysate: sc-126057

## BACKGROUND

Stanniocalcin 1 (STC1) and stanniocalcin 2 (STC2) are mammalian peptide hormones that were previously considered to be present only in bony fish where they are involved in calcium homeostasis. STC1 plays a role in calcium and phosphate homoeostasis and is phosphorylated *in vitro* by protein kinase C, and STC2 is phosphorylated *in vitro* by casein kinase II (CK2). A human fibrosarcoma cell line, HT1080, expresses both STC1 and STC2 as secreted phosphoproteins *in vivo*, with STC2 being phosphorylated by an ecto-CK2-like enzyme. STC1 and STC2 have opposite effects on calcium and phosphate homeostasis, namely anti-hypercalcemic and anti-hypocalcemic actions, respectively. STC1 and STC2 are detected in human adrenal tumors, such as pheochromocytoma, differentiated neuroblastoma aldosterone-producing adenoma, and in cultured adrenal tumor cells (rat pheochromocytoma PC-12 cells and human neuroblastoma NB-1 cells).

## REFERENCES

1. Chang, A.C. and Reddel, R.R. 1998. Identification of a second stanniocalcin cDNA in mouse and human: stanniocalcin 2. Mol. Cell. Endocrinol. 141: 95-99.
2. Honda, S., Kashiwagi, M., Ookata, K., Tojo, A. and Hirose, S. 1999. Regulation by 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> of expression of stanniocalcin messages in the rat kidney and ovary. FEBS Lett. 459: 119-122.
3. Jellinek, D.A., Chang, A.C., Larsen, M.R., Wang, X., Robinson, P.J. and Reddel, R.R. 2000. Stanniocalcin 1 and 2 are secreted as phosphoproteins from human fibrosarcoma cells. Biochem. J. 350: 453-461.
4. Miura, W., Mizunashi, K., Kimura, N., Koide, Y., Noshiro, T., Miura, Y., Furukawa, Y. and Nagura, H. 2000. Expression of stanniocalcin in zona glomerulosa and medulla of normal human adrenal glands, and some adrenal tumors and cell lines. APMIS 108: 367-372.
5. Stasko, S.E. and Wagner, G.F. 2001. Stanniocalcin gene expression during mouse urogenital development: a possible role in mesenchymal-epithelial signalling. Dev. Dyn. 220: 49-59.
6. McCudden, C.R., et al. 2002. Characterization of mammalian stanniocalcin receptors. Mitochondrial targeting of ligand and receptor for regulation of cellular metabolism. J. Biol. Chem. 277: 45249-45258.
7. Yahata, K., et al. 2003. Regulation of stanniocalcin 1 and 2 expression in the kidney by klotho gene. Biochem. Biophys. Res. Commun. 310: 128-134.
8. Ito, D., et al. 2004. Characterization of stanniocalcin 2, a novel target of the mammalian unfolded protein response with cytoprotective properties. Mol. Cell. Biol. 24: 9456-9469.
9. Paciga, M., et al. 2004. Regulation of luteal cell big stanniocalcin production and secretion. Endocrinology 145: 4204-4212.

## CHROMOSOMAL LOCATION

Genetic locus: Stc2 (mouse) mapping to 11 A4.

## PRODUCT

STC2 (m): 293T Lysate represents a lysate of mouse STC2 transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## APPLICATIONS

STC2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive STC2 antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.