



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ADRM1 (m): 293T Lysate: sc-126395

BACKGROUND

ADRM1 (adhesion-regulating molecule 1), also known as GP110, ARM-1 or Rpn13, is a 407 amino acid protein that localizes to both the nucleus and the cytoplasm and is thought to be involved in protein recruitment and cell adhesion. An integral membrane protein, ADRM1 functions to recruit UCH-L5, a deubiquitinating enzyme, to the 26S proteasome, and once at the proteasome it promotes the activity of UCH-L5. Additionally, ADRM1 is thought to mediate lymphocyte adhesion in endothelial cells and may thus play a role in lymphocyte homing. ADRM1 expression is induced by IFN- γ in some cancer cell lines and its expression is upregulated in other metastatic cells, suggesting a role in carcinogenesis. Two isoforms of ADRM1 exist due to alternative splicing events.

REFERENCES

- Shimada, S., Ogawa, M., Takahashi, M., Schlom, J. and Greiner, J.W. 1994. Molecular cloning and characterization of the complementary DNA of an M_r 110,000 antigen expressed by human gastric carcinoma cells and upregulated by γ -interferon. *Cancer Res.* 54: 3831-3836.
- Simins, A.B., Weighardt, H., Weidner, K.M., Weidle, U.H. and Holzmann, B. 1999. Functional cloning of ARM-1, an adhesion-regulating molecule upregulated in metastatic tumor cells. *Clin. Exp. Metastasis* 17: 641-648.
- Lamerant, N. and Kieda, C. 2005. Adhesion properties of adhesion-regulating molecule 1 protein on endothelial cells. *FEBS J.* 272: 1833-1844.
- Hamazaki, J., Iemura, S., Natsume, T., Yashiroda, H., Tanaka, K. and Murata, S. 2006. A novel proteasome interacting protein recruits the deubiquitinating enzyme UCH-37 to 26S proteasomes. *EMBO J.* 25: 4524-4536.
- Qiu, X.B., Ouyang, S.Y., Li, C.J., Miao, S., Wang, L. and Goldberg, A.L. 2006. hRpn13/ADRM1/GP110 is a novel proteasome subunit that binds the deubiquitinating enzyme, UCH-37. *EMBO J.* 25: 5742-5753.
- Cherix, N., Froquet, R., Charette, S.J., Blanc, C., Letourneur, F. and Cosson, P. 2006. A Phg2-ADRM1 pathway participates in the nutrient-controlled developmental response in *Dictyostelium*. *Mol. Biol. Cell* 17: 4982-4987.
- Yao, T., Song, L., Xu, W., DeMartino, G.N., Florens, L., Swanson, S.K., Washburn, M.P., Conaway, R.C., Conaway, J.W. and Cohen, R.E. 2006. Proteasome recruitment and activation of the Uch37 deubiquitinating enzyme by ADRM1. *Nat. Cell Biol.* 8: 994-1002.

CHROMOSOMAL LOCATION

Genetic locus: *Adrm1* (mouse) mapping to 2 H4.

PRODUCT

ADRM1 (m): 293T Lysate represents a lysate of mouse ADRM1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20 $^{\circ}$ C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

ADRM1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ADRM1 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ADRM1 (G-3): sc-374018 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse ADRM1 expression in ADRM1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



ADRM1 (G-3): sc-374018. Western blot analysis of ADRM1 expression in non-transfected: sc-117752 (A) and mouse ADRM1 transfected: sc-126395 (B) 293T whole cell lysates.



ADRM1 (C-7): sc-376221. Western blot analysis of ADRM1 expression in non-transfected: sc-117752 (A) and mouse ADRM1 transfected: sc-126395 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.