



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ARP-1 (m): 293T Lysate: sc-126444

BACKGROUND

COUP (chicken ovalbumin upstream promoter) transcription factors have been cloned in several species and identified as orphan members of the steroid/thyroid hormone receptor superfamily. COUP-TFI (also designated COUP or EAR-3) and ARP-1 (also designated COUP-TFII) exhibit highly regulated and overlapping expression in most tissues. COUP-TFs are highly expressed in the developing and central nervous system, suggesting that these factors may be important in neural development and differentiation. COUP-TFs can compete for binding to response elements which are common to other members of this family, including RAR, RXR, PPAR, HNF-4, VDR and TR. They have been shown to act as negative regulators as well as initiators of transcription.

REFERENCES

- Miyajima, N., et al. 1988. Identification of two novel members of erbA superfamily by molecular cloning: the gene products of the two are highly related to each other. *Nucleic Acids Res.* 16: 11057-11074.
- Wang, L.H., et al. 1989. COUP transcription factor is a member of the steroid receptor superfamily. *Nature* 340: 163-166.
- Ladias, J.A.A., et al. 1991. Regulation of the apolipoprotein AI gene by ARP-1, a novel member of the steroid receptor superfamily. *Science* 251: 561-565.
- Umesono, K., et al. 1991. Direct repeats as selective response elements for the thyroid hormone, retinoic acid and vitamin D3 receptors. *Cell* 65: 1255-1266.
- Cooney, A., et al. 1993. Multiple mechanisms of chicken ovalbumin upstream promoter transcription factor-dependent repression of transactivation by the vitamin D, thyroid hormone, and retinoic acid receptors. *J. Biol. Chem.* 268: 4152-4160.
- Jonk, L.J.C., et al. 1994. Cloning and expression during development of three murine members of the COUP family of nuclear orphan receptors. *Mech. Dev.* 47: 81-97.
- Liu, P.X., et al. 1994. An evolutionary conserved COUP-TF binding element in a neural-specific gene and COUP-TF expression patterns support a major role for COUP-TF in neural development. *Mol. Endocrinol.* 8: 1774-1788.
- Qiu, Y., et al. 1996. Chicken ovalbumin upstream promoter-transcription factors and their regulation. *J. Steroid Biochem. Mol. Biol.* 56: 81-85.

CHROMOSOMAL LOCATION

Genetic locus: Nr2f2 (mouse) mapping to 7 D1.

PRODUCT

ARP-1 (m): 293T Lysate represents a lysate of mouse ARP-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

ARP-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive ARP-1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ARP-1 (E-11): sc-271940 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse ARP-1 expression in ARP-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



ARP-1 (E-11): sc-271940. Western blot analysis of ARP-1 expression in non-transfected: sc-117752 (A) and mouse ARP-1 transfected: sc-126444 (B) 293T whole cell lysates.



ARP-1 (B-2): sc-271265. Western blot analysis of ARP-1 expression in non-transfected: sc-117752 (A) and mouse ARP-1 transfected: sc-126444 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.