



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



B-Myb (m): 293T Lysate: sc-126475

BACKGROUND

The highly leukemogenic avian retrovirus E26 contains two oncogenes, v-Myb and v-Ets, which are expressed together as a fusion protein. The cellular homolog of v-Myb, designated c-Myb, encodes a transcription factor. Deletion or disruption of a negative regulatory domain mapping within the carboxy-terminal domain of c-Myb results in enhanced transactivating capacity and in parallel, leads to activation of its ability to transform hemopoietic cells. c-Myb is expressed preferentially, but not exclusively, in immature hemopoietic cells and its expression decreases as cells differentiate. A second member of the Myb proto-oncogene family, B-Myb, encodes a second sequence-specific DNA binding protein. B-Myb RNA levels are low or undetectable in quiescent cells but increase at the G₁/S-phase transition following mitogenic stimulation. Studies suggest that B-Myb expression rescues cells from p53-induced G₁ arrest mediated by p21.

REFERENCES

1. Gonda, T.J., et al. 1984. Expression of Myb, Myc and Fos proto-oncogenes during the differentiation of a murine myeloid leukaemia. *Nature* 310: 249-251.
2. Gonda, T.J., et al. 1985. Nucleotide sequence of cDNA clones of the murine Myb proto-oncogene. *EMBO J.* 4: 2004-2008.
3. Sakura, H., et al. 1989. Delineation of three functional domains of the transcriptional activator encoded by the c-Myb proto-oncogene. *Proc. Natl. Acad. Sci. USA* 86: 5758-5762.
4. Mizuguchi, G., et al. 1990. DNA binding activity and transcriptional activator function of the human B-Myb protein compared with c-Myb. *J. Biol. Chem.* 265: 9280-9284.
5. Ramsay, R.G., et al. 1991. Increase in specific DNA binding by carboxyl truncation suggests a mechanism for activation of Myb. *Oncogene* 6: 1875-1879.
6. Favier, D., et al. 1994. Detection of proteins that bind to the leucine zipper motif of c-Myb. *Oncogene* 9: 305-311.

CHROMOSOMAL LOCATION

Genetic locus: Mybl2 (mouse) mapping to 2 H2.

PRODUCT

B-Myb (m): 293T Lysate represents a lysate of mouse B-Myb transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

B-Myb (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive B-Myb antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.