



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CAT (m): 293T Lysate: sc-126580

BACKGROUND

Aging affects oxidative metabolism in liver and other tissues. Carnitine acyltransferases are key enzymes of this process in mitochondria. Carnitine acetyltransferase (CAT, CRAT) catalyzes the reversible conversion of acetyl-CoA and carnitine to acetylcarnitine and CoA. The essential functions of CAT are to regenerate CoA, which allows peroxisomal β -oxidation to proceed, and to facilitate transport of acetyl moieties to mitochondria for oxidation. More than 70% of CAT is located in the mitochondrial matrix and it is also located in the endoplasmic reticulum, peroxisomal and mitochondrial inner membrane. An age associated decrease in CAT activity has been reported in many rat systems. The human gene encoding CAT maps to chromosome 9q34.1 and encodes a protein that contains a peroxisomal targeting signal and is expressed mostly in skeletal muscle, and less in heart, liver and pancreas. Total CAT activity is induced by acetate and fatty acids, and repressed by glucose.

REFERENCES

1. Corti, O., Finocchiaro, G., Rossi, E., Zuffardi, O. and DiDonato, S. 1994. Molecular cloning of cDNAs encoding human carnitine acetyltransferase and mapping of the corresponding gene to chromosome 9q34.1. *Genomics* 23: 94-99.
2. Stemple, C.J., Davis, M.A. and Hynes, M.J. 1998. The facC gene of *Aspergillus nidulans* encodes an acetate-inducible carnitine acetyltransferase. *J. Bacteriol.* 180: 6242-6251.
3. Masterson, C. and Wood, C. 2000. Pea chloroplast carnitine acetyltransferase. *Proc. Biol. Sci.* 267: 1-6.
4. Liu, J., Killilea, D.W. and Ames, B.N. 2002. Age-associated mitochondrial oxidative decay: improvement of carnitine acetyltransferase substrate-binding affinity and activity in brain by feeding old rats acetyl-L-carnitine and/or R- α -lipoic acid. *Proc. Natl. Acad. Sci. USA* 99: 1876-1881.
5. SWISS-PROT/TrEMBL (P43155). World Wide Web URL:
<http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: Crat (mouse) mapping to 2 B.

PRODUCT

CAT (m): 293T Lysate represents a lysate of mouse CAT transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

CAT (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive CAT antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.