



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cdc123 (m2): 293T Lysate: sc-126609

BACKGROUND

The eukaryotic cell division cycle consists of a number of gene-controlled sequences that involve cyclin dependent kinases (Cdks) and cell division control (Cdc) proteins. Cdc123 (cell division cycle protein 123), also known as D123, is a 336 amino acid cytoplasmic protein that is involved in cell cycle control. Widely expressed with high expression in thymus, spleen, ovary, testis, small intestine and leukocytes, Cdc123 functions to destabilize Chfr (checkpoint with forkhead and RING finger domain) proteins which, when accumulated, block the G to S phase transition. Cdc123 prevents the Chfr proteins from collecting in the cell, thereby allowing the cell to enter the S phase. Due to its role in cell cycle control, Cdc123 is thought to be a basal marker for luminal breast cancers.

REFERENCES

- Okuda, A. and Kimura, G. 1996. An amino acid change in novel protein D123 is responsible for temperature-sensitive G₁-phase arrest in a mutant of rat fibroblast line 3Y1. *Exp. Cell Res.* 223: 242-249.
- Onisto, M., Zeilante, P., Scannapieco, P., Pellati, D., Pozza, M., Caenazzo, C., Negro, A. and Garbisa, S. 1998. Expression study on D123 gene product: evidence for high positivity in testis. *Exp. Cell Res.* 242: 451-459.
- Okuda, A., Ohtsu, M. and Kimura, G. 1999. Extensive degradation of mutant-type D123 protein is responsible for temperature-sensitive proliferation inhibition in 3Y1tsD123 cells. *Cell Struct. Funct.* 24: 443-449.
- Liu, L.X., Heveker, N., Fackler, O.T., Arold, S., Le Gall, S., Janvier, K., Peterlin, B.M., Dumas, C., Schwartz, O., Benichou, S. and Benarous, R. 2000. Mutation of a conserved residue (D123) required for oligomerization of human immunodeficiency virus type 1 Nef protein abolishes interaction with human thioesterase and results in impairment of Nef biological functions. *J. Virol.* 74: 5310-5319.
- Okuda, A., Ohtsu, M. and Kimura, G. 2001. Reversion of temperature-sensitive mutation by inhibition of proteasome-mediated degradation of mutated D123 protein. *Cell Struct. Funct.* 26: 205-214.
- Biegenowski, P., Shilinski, K., Tschlis, P.N. and Brenner, C. 2004. Cdc123 and checkpoint forkhead associated with RING proteins control the cell cycle by controlling eIF2 γ abundance. *J. Biol. Chem.* 279: 44656-44666.
- Adélaïde, J., Finetti, P., Bekhouche, I., Repellini, L., Geneix, J., Sircoulomb, F., Charafe-Jauffret, E., Cervera, N., Desplans, J., Parzy, D., Schoenmakers, E., Viens, P., Jacquemier, J., Birnbaum, D., Bertucci, F. and Chaffanet, M. 2007. Integrated profiling of basal and luminal breast cancers. *Cancer Res.* 67: 11565-11575.
- Zeggini, E., Scott, L.J., Saxena, R., Voight, B.F., Marchini, J.L., Hu, T., de Bakker, P.I., Abecasis, G.R., Almgren, P., Andersen, G., Ardlie, K., Boström, K.B., Bergman, R.N., Bonnycastle, L.L., Borch-Johnsen, K., et al. 2008. Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat. Genet.* 40: 638-645.

CHROMOSOMAL LOCATION

Genetic locus: Cdc123 (mouse) mapping to 2 A1.

PRODUCT

Cdc123 (m2): 293T Lysate represents a lysate of mouse Cdc123 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Cdc123 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Cdc123 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.