



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# cytohesin-2 (m): 293T Lysate: sc-126697

## BACKGROUND

Cytohesin-2, also known as ARNO (ARF nucleotide-binding-site opener), is an ARF-1 guanine nucleotide exchange factor (GEF). ARF (ADP Ribosylation factor) proteins, a group within the RAS superfamily, are GTP-binding proteins central to the process of vesicle budding. Cytohesin-2 contains an N-terminal coiled-coil domain, a Sec7 domain responsible for GDP/GTP exchange activity, and a C-terminal pleckstrin homology (PH) domain responsible for binding to PIP2. GEF activity of Cytohesin-2 is enhanced by binding of the PH domain to phosphatidylinositol 4,5-bisphosphate which recruits Cytohesin-2 to membranes. Cytohesin-2 is localized to the plasma membrane in mammalian cells and *in vitro* Cytohesin-2 stimulates nucleotide exchange on ARF1 and ARF6.

## REFERENCES

1. Kahn, R.A. and Gilman, A.G. 1984. Purification of a protein cofactor required for ADP-ribosylation of the stimulatory regulatory component of adenylate cyclase by cholera toxin. *J. Biol. Chem.* 259: 6228-6234.
2. Moss, J. and Vaughan, M. 1995. Structure and function of ARF proteins: activators of cholera toxin and critical components of intracellular vesicular transport processes. *J. Biol. Chem.* 270: 12327-12330.
3. Chardin, P., Paris, S., Antonny, B., Robineau, S., Beraud-Dufour, S., Jackson, C.L. and Chabre, M. 1996. A human exchange factor for ARF contains Sec7- and Pleckstrin-homology domains. *Nature* 384: 481-484.
4. Paris, S., Beraud-Dufour, S., Robineau, S., Bigay, J., Antonny, B., Chabre, M. and Chardin, P. 1997. Role of protein-phospholipid interactions in the activation of ARF1 by the guanine nucleotide exchange factor ARNO. *J. Biol. Chem.* 272: 22221-22226.
5. Franco, M., Boretto, J., Sylviane, R., Monier, S., Goud, B., Chardin, P. and Chavrier, P. 1998. ARNO3, a Sec7-domain guanine nucleotide exchange factor for ADP ribosylation factor 1, is involved in the control of Golgi structure and function. *Proc. Natl. Acad. Sci. USA* 95: 9929-9931.
6. Frank, S., Upender, S., Hansen, S. H. and Casanova, J.E. 1998. ARNO is a guanine nucleotide exchange factor for ADP-ribosylation factor 6. *J. Biol. Chem.* 273: 23-27.

## CHROMOSOMAL LOCATION

Genetic locus: Pscd2 (mouse) mapping to 7 B4.

## PRODUCT

cytohesin-2 (m): 293T Lysate represents a lysate of mouse cytohesin-2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

cytohesin-2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive cytohesin-2 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

cytohesin-2 (H-7): sc-374640 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse cytohesin-2 expression in cytohesin-2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



cytohesin-2 (H-7): sc-374640. Western blot analysis of cytohesin-2 expression in non-transfected: sc-117752 (A) and mouse cytohesin-2 transfected: sc-126697 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.