



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FADD (m2): 293T Lysate: sc-126822

BACKGROUND

In contrast to growth factors which promote cell proliferation, FAS ligand (FAS-L) and the tumor necrosis factors (TNFs) rapidly induce apoptosis. Cellular response to FAS-L and TNF is mediated by structurally related receptors containing a conserved "death domain" and belonging to the TNF receptor superfamily. TRADD, FADD and RIP are FAS/TNF-R1 interacting proteins that contain a death domain-homologous region (DDH). TRADD (TNF-R1-associated death domain) and FADD (FAS-associated death domain) associate with the death domains of both FAS and TNF-R1 via their DDH regions. Overexpression of TRADD leads to NF κ B activation and apoptosis in the absence of TNF. Overexpression of FADD causes apoptosis, which can be blocked by the cow pox protein CrmA, suggesting that FADD lies upstream of ICE and possibly other serine proteases. The receptor-interacting protein, RIP, associates with FAS exclusively via its DDH, and this association is abrogated in *lpr* mutants. Unlike TRADD and FADD, RIP contains a putative amino-terminal kinase domain.

REFERENCES

1. Smith, C.A., et al. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation and death. *Cell* 76: 959-962.
2. Nagata, S., et al. 1995. The FAS death factor. *Science* 267: 1449-1456.
3. Sato, T., et al. 1995. FAP-1: a protein tyrosine phosphatase that associates with FAS. *Science* 268: 411-414.
4. Hsu, H., et al. 1995. The TNF receptor 1-associated protein TRADD signals cell death and NF κ B activation. *Cell* 81: 495-504.
5. Baker, S.J., et al. 1996. Transducers of life and death: TNF receptor superfamily and associated proteins. *Oncogene* 12: 1-9.
6. Schneider-Brachert, W., et al. 2006. Inhibition of TNF receptor 1 internalization by adenovirus 14.7K as a novel immune escape mechanism. *J. Clin. Invest.* 116: 2901-2913.
7. Balestrieri, E., et al. 2007. Apoptosis-associated gene expression in HIV-infected patients in response to successful antiretroviral therapy. *J. Med. Virol.* 79: 111-117.
8. Takahashi, R., et al. 2007. Focal adhesion kinase determines the fate of death or survival of cells in response to TNF α in the presence of actinomycin D. *Biochim. Biophys. Acta* 1770: 518-526.
9. Zhu, L., et al. 2007. Decreased expressions of the TNF α signaling adapters in peripheral blood mononuclear cells (PBMCs) are correlated with disease activity in patients with systemic lupus erythematosus. *Clin. Rheumatol.* 26: 1481-1489.

CHROMOSOMAL LOCATION

Genetic locus: *Fadd* (mouse) mapping to 7 F5.

PRODUCT

FADD (m2): 293T Lysate represents a lysate of mouse FADD transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

FADD (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive FADD antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

FADD (G-4): sc-271748 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse FADD expression in FADD transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



FADD (G-4): sc-271748. Western blot analysis of FADD expression in non-transfected: sc-117752 (A) and mouse FADD transfected: sc-126822 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.