



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

# HNF-4 $\alpha$ (m): 293T Lysate: sc-126960

## BACKGROUND

HNF-1 ( $\alpha$  and  $\beta$ ), HNF-3 ( $\alpha$ ,  $\beta$  and  $\gamma$ ), HNF-4 ( $\alpha$  and  $\gamma$ ) and HNF-6 compose, in part, a homeoprotein family designated the hepatocyte nuclear factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues, such as kidney, small intestine and thymus. HNF-3 $\alpha$ , HNF-3 $\beta$  and HNF-3 $\gamma$  regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3 $\alpha$  and HNF-3 $\beta$  have also been shown to be involved in gastrulation events such as body axis formation. HNF-4 $\alpha$  and HNF-4 $\gamma$  have been shown to be important for early embryo development. HNF-4 $\alpha$  is expressed in liver, kidney, pancreas, small intestine, testis and colon, while HNF-4 $\gamma$  is expressed in each of these tissues except liver. HNF-6 has been shown to bind to the promoter of HNF-3 $\beta$ , which indicates a potential role of HNF-6 in gut endoderm epithelial cell differentiation. Evidence suggests that HNF-6 may also be a transcriptional activator for at least 22 other hepatocyte-enriched genes, including cytochrome P450 2C13 and  $\alpha$ -1 antitrypsin.

## REFERENCES

1. Bach, I., et al. 1993. More potent transcriptional activators or a trans-dominant inhibitor of the HNF-1 homeoprotein family are generated by alternative RNA processing. *EMBO J.* 12: 4229-4242.
2. Kaestner, K.H., et al. 1994. The HNF-3 gene family of transcription factors in mice: gene structure, cDNA sequence, and mRNA distribution. *Genomics* 20: 377-385.
3. Drewes, T., et al. 1996. Human hepatocyte nuclear factor-4 isoforms are encoded by distinct and differentially expressed genes. *Mol. Cell. Biol.* 16: 925-931.
4. Samadani, U., et al. 1996. The transcriptional activator hepatocyte nuclear factor-6 regulates liver gene expression. *Mol. Cell. Biol.* 16: 6273-6284.
5. Kojima, K., et al. 2006. The expression of hepatocyte nuclear factor-4 $\alpha$ , a developmental regulator of visceral endoderm, correlates with the intestinal phenotype of gastric adenocarcinomas. *Pathology* 38: 548-554.
6. Gupta, R.K., et al. 2007. Expansion of adult  $\beta$  cell mass in response to increased metabolic demand is dependent on HNF-4 $\alpha$ . *Genes Dev.* 21: 756-769.
7. Krajewski, J., et al. 2007. Interleukin-1 $\beta$  inhibits the hypoxic inducibility of the erythropoietin enhancer by suppressing hepatocyte nuclear factor-4 $\alpha$ . *Cell. Mol. Life Sci.* 64: 989-998.
8. Ogasawara, K., et al. 2007. Hepatocyte nuclear factor-4 $\alpha$  regulates the human organic anion transporter 1 gene in the kidney. *Am. J. Physiol. Renal Physiol.* 292: F1819-1826.
9. Klapper, M., et al. 2007. The human intestinal fatty acid binding protein (hFABP2) gene is regulated by HNF-4 $\alpha$ . *Biochem. Biophys. Res. Commun.* 356: 147-152.

## CHROMOSOMAL LOCATION

Genetic locus: Hnf4a (mouse) mapping to 2 H3.

## PRODUCT

HNF-4 $\alpha$  (m): 293T Lysate represents a lysate of mouse HNF-4 $\alpha$  transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## APPLICATIONS

HNF-4 $\alpha$  (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive HNF-4 $\alpha$  antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20 $^{\circ}$  C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.