



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



IL-4R α (m): 293T Lysate: sc-127006

BACKGROUND

The IL-2 receptor is a multicomponent complex consisting of three subunits, α , β and γ , each of which is required for high affinity binding of IL-2. The α chain functions primarily in binding IL-2, whereas the β and γ chains contribute to IL-2 binding and are essential to IL-2-induced activation of signaling pathways leading to T cell growth. Both IL-4R and IL-7R were initially described as single chain, high-affinity ligand-binding cytokine receptors. However, it is now well established that the IL-2R γ chain functions as a second subunit of the high affinity IL-4R and IL-7R receptors. Consequently, the originally described subunits of these latter receptors are now referred to as IL-4R α and IL-7R α , respectively, while the common subunit is referred to as γ c. Although the common γ chain enhances ligand binding in these three cytokine receptors, it has no capacity to bind these ligands on its own. There is evidence that the γ c chain is also a subunit of IL-13R.

REFERENCES

1. Mosley, B., Beckmann, M.P., March, C.J., Idzerda, R.L., Gimpel, S.D., VandenBos, T., Friend, D., Alpert, A., Anderson, D., Jackson, J., Wignall, J.M., Smith, C., Gallis, B., Sims, J.E., Urdal, D., Widmer, M.B., et al. 1989. The murine interleukin-4 receptor: molecular cloning and characterization of secreted and membrane bound forms. *Cell* 59: 335-348.
2. Goodwin, R.G., Friend, D., Ziegler, S.F., Jerzy, R., Falk, B.A., Gimpel, S., Cosman, D., Dower, S.K., March, C.J., Namen, A.E. and Park, L.S. 1990. Cloning of the human and murine interleukin-7 receptors: demonstration of a soluble form and homology to a new receptor superfamily. *Cell* 60: 941-951.
3. Takeshita, T., Asao, H., Ohtani, K., Ishii, N., Kumaki, S., Tanaka, N., Munakata, H., Nakamura, M. and Sugamura, K. 1992. Cloning of the γ chain of the human IL-2 receptor. *Science* 257: 379-382.
4. Cao, X., Kozak, C.A., Liu, Y.J., Noguchi, M., O'Connell, E. and Leonard, W.J. 1993. Characterization of cDNAs encoding the murine interleukin 2 receptor (IL-2R) γ chain: chromosomal mapping and tissue specificity of IL-2R γ chain expression. *Proc. Natl. Acad. Sci. USA* 90: 8464-8468.
5. Kondo, M., Takeshita, T., Ishii, N., Nakamura, M., Watanabe, S., Arai, K.I. and Sugamura, K. 1993. Sharing of the interleukin-2 (IL-2) receptor γ chain between receptors for IL-2 and IL-4. *Science* 262: 1874-1877.
6. Russell, S.M., Keegan, A.D., Harada, N., Nakamura, Y., Noguchi, M., Leland, P., Friedmann, M.C., Miyajima, A., Puri, R.K., Paul, W.E. and Leonard, W.J. 1993. Interleukin-2 receptor γ chain: a functional component of the interleukin-4 receptor. *Science* 262: 1880-1883.
7. Minami, Y., Kono, T., Miyazaki, T. and Taniguchi, T. 1993. The IL-2 receptor complex: its structure, function and target genes. *Annu. Rev. Immunol.* 11: 245-268.
8. He, Y.W., Adkins, B., Furse, R.K. and Malek, T.R. 1995. Expression and function of the γ c subunit of the IL-2, IL-4 and IL-7 receptors. *J. Immunol.* 154: 1596-1605.

CHROMOSOMAL LOCATION

Genetic locus: Il4r (mouse) mapping to 7 F3.

PRODUCT

IL-4R α (m): 293T Lysate represents a lysate of mouse IL-4R α transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

IL-4R α (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive IL-4R α antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


IL-4R α (D-2): sc-137257 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse IL-4R α expression in IL-4R α transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



IL-4R α (D-2): sc-137257. Western blot analysis of IL-4R α expression in non-transfected: sc-117752 (**A**) and truncated mouse IL-4R α transfected: sc-127006 (**B**) 293T whole cell lysates.

IL-4R α (G-9): sc-137256. Western blot analysis of IL-4R α expression in non-transfected: sc-117752 (**A**) and truncated mouse IL-4R α transfected: sc-127006 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.