



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# IP3KA (m): 293T Lysate: sc-127014

## BACKGROUND

Inositol 1,4,5-trisphosphate (Ins(1,4,5)P<sub>3</sub>) regulates the level of calcium within the cell by releasing calcium from intracellular stores. Ins(1,4,5)P<sub>3</sub> is phosphorylated by inositol 1,4,5-trisphosphate 3-kinase (IP3K) to form inositol 1,3,4,5-tetrakisphosphate (Ins(1,4,5)P<sub>4</sub>), which is thought to regulate the influx of calcium across the plasma membrane. IP3K exists as three isoforms, IP3KA, B, and C. IP3KA, the most highly characterized isoform, is expressed in rat brain and testis. IP3KB is expressed in various rat tissues such as lung, thymus, testis, brain, and heart. IP3K activity is stimulated in the presence of calmodulin via phosphorylation by cAMP-dependent protein kinase, protein kinase C, or calcium/calmodulin dependent protein kinase II and, subsequently, mediates the inositol phosphate signaling pathways.

## REFERENCES

1. Johanson, R.A., Hansen, C.A. and Williamson, J.R. 1988. Purification of D-myo-inositol 1,4,5-trisphosphate 3-kinase from rat brain. *J. Biol. Chem.* 263: 7465-7471.
2. Berridge, M.J. and Irvine, R.F. 1989. Inositol phosphates and cell signaling. *Nature* 341: 197-205.
3. Sim, S.S., Kim, J.W. and Rhee, S.G. 1990. Regulation of D-myo-inositol 1,4,5-trisphosphate 3-kinase by cAMP-dependent protein kinase and protein kinase C. *J. Biol. Chem.* 265: 10367-10372.
4. Takazawa, K., Vandekerckhove, J., Dumont, J.E. and Erneux, C. 1990. Cloning and expression in *Escherichia coli* of a rat brain cDNA encoding a Ca<sup>2+</sup>/calmodulin-sensitive inositol 1,4,5-trisphosphate 3-kinase. *Biochem. J.* 272: 107-112.
5. Irvine, R.F. 1991. Inositol tetrakisphosphate as a second messenger: confusions, contradictions, and a potential resolution. *Bioessays* 13: 419-427.
6. Vanweyenberg, V., Communi, D., D'Santos, C.S. and Erneux, C. 1995. Tissue and cell-specific expression of Ins(1,4,5)P<sub>3</sub> 3-kinase isoenzymes. *Biochem. J.* 306: 429-435.
7. Woodring, P.J. and Garrison, J.C. 1997. Expression, purification, and regulation of two isoforms of the inositol 1,4,5-trisphosphate 3-kinase. *J. Biol. Chem.* 272: 30447-30454.

## CHROMOSOMAL LOCATION

Genetic locus: *Itpka* (mouse) mapping to 2 E5.

## PRODUCT

IP3KA (m): 293T Lysate represents a lysate of mouse IP3KA transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

IP3KA (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive IP3KA antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


IP3KA (F-3): sc-271838 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse IP3KA expression in IP3KA transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG<sub>κ</sub> BP-HRP: sc-516102 or m-IgG<sub>κ</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



IP3KA (F-3): sc-271838. Western blot analysis of IP3KA expression in non-transfected: sc-117752 (**A**) and mouse IP3KA transfected: sc-127014 (**B**) 293T whole cell lysates.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.