



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Nrf2 (m): 293T Lysate: sc-127242

BACKGROUND

The NF-E2 DNA binding protein is composed of two subunits, p45 and MafK. It regulates expression of globin genes in developing erythroid cells through interaction with Maf recognition elements (Mares). A family of NF-E2-related proteins, which are collectively known as the Cap 'n' collar (CNC) family and include Nrf1 (also designated TCF11), Nrf2 and Nrf3, are bZIP transcription factors that heterodimerize with Maf proteins to bind Maf sequences. The Nrf proteins also bind the antioxidant response element (ARE) and are implicated in the regulation of detoxification enzymes and the oxidative stress response. They do so by heterodimerizing with Jun family members (c-Jun, Jun B and Jun D) to activate gene expression, specifically the detoxifying enzyme NQO1. The genes encoding Nrf1 and Nrf2 map to human chromosomes 17q21.32 and 2q31.2, respectively. Nrf2 is widely expressed and is thought to translocate to the nucleus after treatment with xenobiotics and antioxidants, which stimulate its release from its repressor protein, Keap1. The gene encoding human Nrf3 maps to chromosome 7p15.2. Nrf3 is highly expressed in placenta, B cells and monocytes.

REFERENCES

- Chan, J.Y., et al. 1995. Chromosomal localization of the human NF-E2 family of bZIP transcription factors by fluorescence *in situ* hybridization. *Hum. Genet.* 95: 265-269.
- Chan, K., et al. 1996. Nrf2, a member of the NF-E2 family of transcription factors, is not essential for murine erythropoiesis, growth, and development. *Proc. Natl. Acad. Sci. USA* 93: 13943-13948.
- Kobayashi, A., et al. 1999. Molecular cloning and functional characterization of a new Cap 'n' collar family transcription factor Nrf3. *J. Biol. Chem.* 274: 6443-6452.
- Chan, J.Y., et al. 2000. Impaired expression of glutathione synthetic enzyme genes in mice with targeted deletion of the Nrf2 basic-leucine zipper protein. *Biochim. Biophys. Acta* 1517: 19-26.
- Dhakshinamoorthy, S., et al. 2000. Small Maf (MafG and MafK) proteins negatively regulate antioxidant response element-mediated expression and antioxidant induction of the NAD(P)H: Quinone oxidoreductase1 gene. *J. Biol. Chem.* 275: 40134-40141.
- Jaiswal, A.K. 2000. Regulation of genes encoding NAD(P)H: quinone oxidoreductases. *Free Radic. Biol. Med.* 29: 254-262.
- Myhrstad, M.C., et al. 2001. TCF11/Nrf1 overexpression increases the intracellular glutathione level and can transactivate the γ -glutamylcysteine synthetase (GCS) heavy subunit promoter. *Biochim. Biophys. Acta* 1517: 212-219.
- Jakel, R.J., et al. 2007. Nrf2-mediated protection against 6-hydroxydopamine. *Brain Res.* 1144: 192-201.
- Marzec, J.M., et al. 2007. Functional polymorphisms in the transcription factor Nrf2 in humans increase the risk of acute lung injury. *FASEB J.* 21: 2237-2246.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

CHROMOSOMAL LOCATION

Genetic locus: Nfe2l2 (mouse) mapping to 2 C3.

PRODUCT

Nrf2 (m): 293T Lysate represents a lysate of mouse Nrf2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Nrf2 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Nrf2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.