



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# PGRP-S (m): 293T Lysate: sc-127321

## BACKGROUND

Peptidoglycan recognition proteins (PGRPs) are molecules that recognize peptidoglycan, a large component in bacterial cell walls. In insects, PGRPs activate antimicrobial pathways, and in mammals PGRPs function as anti-bacterial neutrophil proteins. PGRP-L halts bacterial growth by acting as an alanine amidase, an enzyme that hydrolyzes the amide bond of bacterial peptidoglycan. PGRP- $\alpha$  and PGRP- $\beta$  are also members of the PGRP family that help recognize bacteria by binding to peptidoglycan and Gram Positive Bacteria, but they do not have amidase activity. PGRP-S participates in intracellular killing of Gram Positive Bacteria by stimulating two antimicrobial defense systems, the prophenoloxidase cascade and the antimicrobial peptides through toll receptors.

## REFERENCES

- Liu, C., Xu, Z., Gupta, D. and Dziarski, R. 2001. Peptidoglycan recognition proteins: a novel family of four human innate immunity pattern recognition molecules. *J. Biol. Chem.* 276: 34686-34694.
- Dziarski, R., Platt, K.A., Gelius, E., Steiner, H. and Gupta, D. 2003. Defect in neutrophil killing and increased susceptibility to infection with non-pathogenic Gram Positive Bacteria in peptidoglycan recognition protein-S (PGRP-S)-deficient mice. *Blood* 102: 689-697.
- Kibardin, A.V., Mirkina, I.I., Zakeeva, I.R., Baranova, E.V., Georgiev, G.P. and Kiselev, S.L. 2003. Expression analysis of proteins encoded by genes of the TAG7/TAGL (PGRP-S,-L) family in human peripheral blood cells. *Genetika* 39: 244-249.
- Lo, D., Tynan, W., Dickerson, J., Mendy, J., Chang, H.W., Scharf, M., Byrne, D., Brayden, D., Higgins, L., Evans, C. and O'Mahony, D.J. 2003. Peptidoglycan recognition protein expression in mouse Peyer's Patch follicle associated epithelium suggests functional specialization. *Cell. Immunol.* 224: 8-16.
- Wang, Z.M., Li, X., Cocklin, R.R., Wang, M., Wang, M., Fukase, K., Inamura, S., Kusumoto, S., Gupta, D. and Dziarski, R. 2003. Human peptidoglycan recognition protein-L is an N-acetylmuramoyl-L-alanine amidase. *J. Biol. Chem.* 278: 49044-49052.
- Cho, J.H., Fraser, I.P., Fukase, K., Kusumoto, S., Fujimoto, Y., Stahl, G.L. and Ezekowitz, R.A. 2005. Human peptidoglycan recognition protein S is an effector of neutrophil-mediated innate immunity. *Blood* 106: 2551-2558.
- Guan, R., Wang, Q., Sundberg, E.J. and Mariuzza, R.A. 2005. Crystal structure of human peptidoglycan recognition protein S (PGRP-S) at 1.70 Å resolution. *J. Mol. Biol.* 347: 683-691.
- Kumar, S., Roychowdhury, A., Ember, B., Wang, Q., Guan, R., Mariuzza, R.A. and Booms, G.J. 2005. Selective recognition of synthetic lysine and meso-diaminopimelic acid-type peptidoglycan fragments by human peptidoglycan recognition proteins- $\alpha$  and -S. *J. Biol. Chem.* 280: 37005-37012.
- Tydell, C.C., Yuan, J., Tran, P. and Selsted, M.E. 2006. Bovine peptidoglycan recognition protein-S: antimicrobial activity, localization, secretion, and binding properties. *J. Immunol.* 176: 1154-1162.

## CHROMOSOMAL LOCATION

Genetic locus: Pglyrp1 (mouse) mapping to 7 A3.

## PRODUCT

PGRP-S (m): 293T Lysate represents a lysate of mouse PGRP-S transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

PGRP-S (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PGRP-S antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PGRP-S (G-1): sc-365304 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse PGRP-S expression in PGRP-S transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



PGRP-S (G-1): sc-365304. Western blot analysis of PGRP-S expression in non-transfected: sc-117752 (**A**) and mouse PGRP-S transfected: sc-127321 (**B**) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.