



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PME-1 (m2): 293T Lysate: sc-127356

BACKGROUND

Protein phosphatase methylesterase-1 (PME-1) catalyzes the demethylation and inactivation of protein phosphatase (PP2A), which is a multimeric phosphoserine/threonine protein phosphatase associated with growth inhibition and cell cycle arrest. Carboxymethylation and demethylation is a covalent modification that regulates the catalytic activity of certain proteins in eukaryotes. Electrostatic interactions that occur at residues or metals in or near the active site can influence the specificity of carboxymethylation and demethylation. PME-1 can demethylate PP2A catalytic subunit *in vitro* and okadaic acid treatment is capable of inhibiting this reaction. PME-1 is conserved from yeast to human and contains a motif found in lipases having a catalytic triad-activated serine as their active site nucleophile.

REFERENCES

- Lee, J., Chen, Y., Tolstykh, T. and Stock, J. 1996. A specific protein carboxyl methyltransferase that demethylates phosphoprotein phosphatase 2A in bovine brain. *Proc. Natl. Acad. Sci. USA* 93: 6043-6047.
- Schonthal, A.H. 1998. Role of PP2A in intracellular signal transduction pathways. *Front. Biosci.* 3: D1262-D1273.
- Ogris, E., Du, X., Nelson, K.C., Mak, E.K., Yu, X.X., Lane, W.S. and Pallas, D.C. 1999. A protein phosphatase methylesterase (PME-1) is one of several novel proteins stably associating with two inactive mutants of protein phosphatase 2A. *J. Biol. Chem.* 274: 14382-14391.
- Wu, J., Tolstykh, T., Lee, J., Boyd, K., Stock, J.B. and Broach, J.R. 2000. Carboxyl methylation of the phosphoprotein phosphatase 2A catalytic subunit promotes its functional association with regulatory subunits *in vivo*. *EMBO J.* 19: 5672-5681.
- Tolstykh, T., Lee, J., Vafai, S. and Stock, J.B. 2000. Carboxyl methylation regulates phosphoprotein phosphatase 2A by controlling the association of regulatory B subunits. *EMBO J.* 19: 5682-5691.
- Gagnon, S.N., Hengartner, M.O. and Desnoyers, S. 2002. The genes PME-1 and PME-2 encode two poly(ADP-ribose) polymerases in *Caenorhabditis elegans*. *Biochem. J.* 368: 263-271.
- Longin, S., Jordens, J., Martens, E., Stevens, I., Janssens, V., Rondelez, E., De Baere, I., Derua, R., Waelkens, E., Goris, J. and Van Hoof, C. 2004. An inactive protein phosphatase 2A population is associated with methyltransferase and can be re-activated by the phosphotyrosyl phosphatase activator. *Biochem. J.* 380: 111-119.

CHROMOSOMAL LOCATION

Genetic locus: Ppme1 (mouse) mapping to 7 E3.

PRODUCT

PME-1 (m2): 293T Lysate represents a lysate of mouse PME-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

PME-1 (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PME-1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PME-1 (A-10): sc-137145 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse PME-1 expression in PME-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PME-1 (A-10): sc-137145. Western blot analysis of PME-1 expression in non-transfected: sc-117752 (A) and mouse PME-1 transfected: sc-127356 (B) 293T whole cell lysates.



PME-1 (B-12): sc-25278. Western blot analysis of PME-1 expression in non-transfected: sc-117752 (A) and mouse PME-1 transfected: sc-127356 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.