



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Wee 1 (m): 293T Lysate: sc-127786

## BACKGROUND

Phosphorylation of Cdc2 on Threonine 14 and Tyrosine 15 is required to maintain Cdc2 in an inactive state throughout the S and G<sub>2</sub> phases of the cell cycle. The human Wee 1 protein, WEE1Hu, encodes a tyrosine-specific protein kinase that phosphorylates Cdc2 on tyrosine 15. Myt 1, a member of the Wee 1 family of protein kinases, has been shown to phosphorylate Cdc2 on both Threonine 14 and Tyrosine 15 in a cyclin-dependent manner. Activity of both Wee 1 Hu and Myt 1 is regulated during the cell cycle, suggesting that both proteins play a role in mitotic control. Dephosphorylation of Cdc2 on Threonine 14 and Tyrosine 15 in late G<sub>2</sub> by Cdc25 then activates the Cdc2/cyclin B complex to allow entry into mitosis.

## REFERENCES

- Morla, A., et al. 1989. Reversible tyrosine phosphorylation of Cdc2: dephosphorylation accompanies activation during entry into mitosis. *Cell* 58: 193-203.
- Krek, W., et al. 1991. Differential phosphorylation of vertebrate p34<sup>Cdc2</sup> kinase at the G<sub>1</sub>/S and G<sub>2</sub>/M transitions of the cell cycle: identification of major phosphorylation sites. *EMBO J.* 10: 305-316.
- Igarashi, M., et al. 1991. Wee 1-like gene in human cells. *Nature* 353: 80-83.
- McGowan, C.H., et al. 1995. Human Wee 1 kinase inhibits cell division by phosphorylating p34<sup>Cdc2</sup> exclusively on Tyr 15. *EMBO J.* 12: 75-85.
- Watanabe, N., et al. 1995. Regulation of the human WEE1Hu Cdk Tyrosine 15 kinase during the cell cycle. *EMBO J.* 14: 1878-1891.
- Liu, F., et al. 1997. The human Myt 1 kinase preferentially phosphorylates Cdc2 on Threonine 14 and localizes to the endoplasmic reticulum and Golgi complex. *Mol. Cell. Biol.* 17: 571-583.
- Squire, C.J., et al. 2005. Structure and inhibition of the human cell cycle checkpoint kinase, Wee 1A kinase: an atypical tyrosine kinase with a key role in Cdk1 regulation. *Structure* 13: 541-550.
- Kiviharju-af Hällström, T.M., et al. 2007. Human prostate epithelium lacks Wee 1A-mediated DNA damage-induced checkpoint enforcement. *Proc. Natl. Acad. Sci. USA* 104: 7211-7216.
- Kamata, M., et al. 2008. Human immunodeficiency virus type 1 Vpr binds to the N lobe of the Wee 1 kinase domain and enhances kinase activity for Cdc2. *J. Virol.* 82: 5672-5682.

## CHROMOSOMAL LOCATION

Genetic locus: Wee1 (mouse) mapping to 7 F1.

## PRODUCT

Wee 1 (m): 293T Lysate represents a lysate of mouse Wee 1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

Wee 1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive Wee 1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Wee 1 (B-11): sc-5285 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse Wee 1 expression in Wee 1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Wee 1 (B-11): sc-5285. Western blot analysis of Wee 1 expression in non-transfected: sc-117752 (A) and mouse Wee 1 transfected: sc-127786 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.