



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Brn-3a (h): 293T Lysate: sc-128117

BACKGROUND

The Brn family of transcription factors are found in a highly restricted subset of neurons and are critical to the early embryonic development of the central nervous system. Brn-1 and Brn-2 are class III POU (Pit-Oct-Unc) domain proteins, whereas Brn-3 is a class IV POU domain protein. Three Brn-3 proteins have been described and are designated Brn-3a, Brn-3b and Brn-3c. While Brn-3a and Brn-3c stimulate transcription, Brn-3b generally functions as a transcriptional repressor. However, Brn-3b, but not Brn-3a, has been shown to regulate the expression of the acetylcholine receptor. Interestingly, Brn-3a has two functional transactivating domains, one at the amino-terminus and one at the carboxy-terminus. Brn-2 is thought to be involved in smooth muscle cell development and differentiation.

REFERENCES

- Collum, R.G., et al. 1992. A novel POU homeodomain gene specifically expressed in cells of the developing mammalian nervous system. *Nucleic Acids Res.* 20: 4919-4925.
- Xiang, M., et al. 1993. The gene for Brn-3b: a POU-domain protein expressed in retinal ganglion cells is assigned to the q31.2 region of chromosome 4. *Human Genome Mapping Workshop* 93: 7.
- Fedtsova, N.G., et al. 1995. Brn-3.0 expression identifies early post-mitotic CNS neurons and sensory neural precursors. *Mech. Dev.* 53: 291-304.
- Schonemann, M.D., et al. 1995. Development and survival of the endocrine hypothalamus and posterior pituitary gland requires the neuronal POU domain factor Brn-2. *Genes Dev.* 9: 3122-3135.
- Budhram-Mahadeo, V., et al. 1996. The different activities of the two activation domains of the Brn-3a transcription factor are dependent on the context of the binding site. *J. Biol. Chem.* 271: 9108-9113.
- Dawson, S.J., et al. 1996. A single amino acid change converts an inhibitory transcription factor into an activator. *J. Biol. Chem.* 271: 11631-11633.
- Erkman, L., et al. 1996. Role of transcription factors Brn-3.1 and Brn-3.2 in auditory and visual system development. *Nature* 381: 603-606.
- Gan, L., et al. 1996. POU domain factor Brn-3b is required for the development of a large set of retinal ganglion cells. *Proc. Natl. Acad. Sci. USA* 93: 3920-3925.
- Suzuki, T., et al. 1996. Preferential differentiation of P19 mouse embryonal carcinoma cells into smooth muscle cells. Use of retinoic acid and antisense against the central nervous system-specific POU transcription factor Brn-2. *Circ. Res.* 78: 395-404.

CHROMOSOMAL LOCATION

Genetic locus: POU4F1 (human) mapping to 13q31.1.

PRODUCT

Brn-3a (h): 293T Lysate represents a lysate of human Brn-3a transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Brn-3a (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Brn-3a antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Brn-3a (14A6): sc-8429 is recommended as a positive control antibody for Western Blot analysis of enhanced human Brn-3a expression in Brn-3a transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Brn-3a (14A6): sc-8429. Near-infrared western blot analysis of Brn-3a expression in non-transfected: sc-117752 (A) and human Brn-3a transfected: sc-128117 (B) 293T whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGκ BP-CFL 680: sc-516180.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.