



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Crk-L (h): 293T Lysate: sc-128367

BACKGROUND

SH2 and SH3 (Src homology) domains were originally identified as critical functional domains within non-receptor proteins with tyrosine kinase activity. A subset of these proteins appears to exist predominately of SH2/SH3 domains in the absence of detectable catalytic domains. One of the first members of the family to be identified, Crk, is a transformation-specific protein that induces elevation of cellular phosphotyrosine levels, but lacks tyrosine kinase activity itself. A second protein, Nck, consists solely of three SH3 domains and one SH2 domain, while GRB2 contains an SH2 domain flanked on both sides by SH3 domains. A member of this protein class, Crk-L, is encoded by a gene located on chromosome 22, band 11, centro-meric of the chronic myelogenous leukemia breakpoint region. Crk-L encodes a 303 amino acid protein with one SH2 and 2 SH3 domains.

REFERENCES

1. Mayer, B., et al. 1988. A novel viral oncogene with structural similarity to phospholipase C. *Nature* 332: 272-275.
2. Lehmann, J.M., et al. 1990. Nck, a melanoma cDNA encoding a cytoplasmic protein consisting of the Src homology units SH2 and SH3. *Nucleic Acids Res.* 18: 1048.
3. Koch, C.A., et al. 1991. SH2 and SH3 domains: elements that control interactions of cytoplasmic signaling proteins. *Science* 252: 668-674.
4. Pawson, T., et al. 1992. SH2 and SH3 domains: from structure to function. *Cell* 71: 359-362.
5. Williams, L.T. 1992. Missing links between receptors and Ras. *Curr. Biol.* 2: 601-603.
6. Park, D., et al. 1992. Phosphorylation of Nck in response to a variety of receptors, phorbol myristate acetate, and cyclic AMP. *Mol. Cell. Biol.* 12: 5816-5823.
7. Lowenstein, E.J., et al. 1992. The SH2 and SH3 domain-containing protein GRB2 links receptor tyrosine kinases to Ras signaling. *Cell* 70: 431-442.

CHROMOSOMAL LOCATION

Genetic locus: CRKL (human) mapping to 22q11.21.

PRODUCT

Crk-L (h): 293T Lysate represents a lysate of human Crk-L transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Crk-L (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Crk-L antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Crk-L (B-1): sc-365092 is recommended as a positive control antibody for Western Blot analysis of enhanced human Crk-L expression in Crk-L transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Crk-L (B-1): sc-365092. Western blot analysis of Crk-L expression in non-transfected: sc-117752 (A) and human Crk-L transfected: sc-128367 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.