



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# GRINL1A (h): 293T Lysate: sc-128737

## BACKGROUND

GRINL1A (glutamate receptor-like protein 1A), also known as Gcom2 or Gdown, is a 148 amino acid protein belonging to the GRINL1 family. The gene encoding GRINL1A maps to human chromosome 15q21.3, and exists as two readthrough transcript variations. Alternative splicing events additionally result in six isoforms, designated Gdown1, Gdown6, isoform 3, Gdown4, Gdown3 and Gcom1-which exists as a naturally occurring fusion protein with GRINL1A. Isoform 1 localizes to the nucleus and is expressed in adult and fetal brain, as well as heart, kidney, skeletal muscle, small intestine, lung, prostate and testis. A component of the Pol III(G) complex, isoform 1 may also be involved in the Mediator complex-dependent regulation of transcription activation.

## REFERENCES

- Roginski, R.S., Mohan Raj, B.K., Finkernagel, S.W. and Sciorra, L.J. 2001. Assignment of an ionotropic glutamate receptor-like gene (GRINL1A) to human chromosome 15q22.1 by *in situ* hybridization. *Cytogenet. Cell Genet.* 93: 143-144.
- Roginski, R.S., Mohan Raj, B.K., Birditt, B. and Rowen, L. 2004. The human GRINL1A gene defines a complex transcription unit, an unusual form of gene organization in eukaryotes. *Genomics* 84: 265-276.
- Roginski, R.S., Goubaeva, F., Mikami, M., Fried-Cassorla, E., Nair, M.R. and Yang, J. 2008. GRINL1A colocalizes with N-methyl D-aspartate receptor NR1 subunit and reduces N-methyl D-aspartate toxicity. *Neuroreport* 19: 1721-1726.
- Seeger, T.S., Frank, D., Rohr, C., Will, R., Just, S., Grund, C., Lyon, R., Luedde, M., Koegl, M., Sheikh, F., Rottbauer, W., Franke, W.W., Katus, H.A., Olson, E.N. and Frey, N. 2010. Myozap, a novel intercalated disc protein, activates serum response factor-dependent signaling and is required to maintain cardiac function *in vivo*. *Circ. Res.* 106: 880-890.
- García-Mayoral, M.F., Martínez-Moreno, M., Albar, J.P., Rodríguez-Crespo, I. and Bruix, M. 2010. Structural basis for the interaction between dynein light chain 1 and the glutamate channel homolog GRINL1A. *FEBS J.* 277: 2340-2350.
- Huo, L., Wen, W., Wang, R., Kam, C., Xia, J., Feng, W. and Zhang, M. 2011. Cdc42-dependent formation of the ZO-1/MRCK $\beta$  complex at the leading edge controls cell migration. *EMBO J.* 30: 665-678.
- Lettre, G., Palmer, C.D., Young, T., Ejebe, K.G., Allayee, H., Benjamin, E.J., Bennett, F., Bowden, D.W., Chakravarti, A., Dreisbach, A., Farlow, D.N., Folsom, A.R., Fornage, M., Forrester, T., Fox, E., Haiman, C.A., Hartiala, J., Harris, T.B., Hazen, S.L., Heckbert, S.R., Henderson, B.E., Hirschhorn, J.N., et al. 2011. Genome-wide association study of coronary heart disease and its risk factors in 8,090 African Americans: the NHLBI care project. *PLoS Genet.* 7: e1001300.

## CHROMOSOMAL LOCATION

Genetic locus: POLR2M (human) mapping to 15q21.3.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

GRINL1A (h): 293T Lysate represents a lysate of human GRINL1A transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## APPLICATIONS

GRINL1A (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive GRINL1A antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.