



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GSTT2 (h3): 293T Lysate: sc-128756

BACKGROUND

Glutathione (GSH) is a tripeptide antioxidant which reduces disulfide bonds between cytoplasmic proteins. The constitutive enzyme glutathione reductase transforms glutathione into its reduced state which ultimately can provide a measure of cellular toxicity. GSTT2 (glutathione S-transferase θ -2), also known as GST class- θ -2, is a 244 amino acid enzyme with sulfatase activity that functions in conjugating reduced glutathione to hydrophobic electrophiles. GSTT2 exists as a homodimer in the cytoplasm and is expressed in low levels in the liver and the lung. GSTT2 belongs to the GST superfamily and contains both a GST C-terminal and a GST N-terminal domain. The gene encoding GSTT2 exists on human chromosome 22.

REFERENCES

1. Hussey, A.J. and Hayes, J.D. 1992. Characterization of a human class- θ glutathione S-transferase with activity towards 1-menaphthyl sulphate. *Biochem. J.* 286: 929-935.
2. Tan, K.L., Webb, G.C., Baker, R.T. and Board, P.G. 1995. Molecular cloning of a cDNA and chromosomal localization of a human θ -class glutathione S-transferase gene (GSTT2) to chromosome 22. *Genomics* 25: 381-387.
3. Mainwaring, G.W., Williams, S.M., Foster, J.R., Tugwood, J. and Green, T. 1996. The distribution of θ -class glutathione S-transferases in the liver and lung of mouse, rat and human. *Biochem. J.* 318: 297-303.
4. Rossjohn, J., McKinstry, W.J., Oakley, A.J., Verger, D., Flanagan, J., Chelvanayagam, G., Tan, K.L., Board, P.G. and Parker, M.W. 1998. Human θ class glutathione transferase: the crystal structure reveals a sulfate-binding pocket within a buried active site. *Structure* 6: 309-322.
5. Sprenger, R., Schlagenhauer, R., Kerb, R., Bruhn, C., Brockmüller, J., Roots, I. and Brinkmann, U. 2000. Characterization of the glutathione S-transferase GSTT1 deletion: discrimination of all genotypes by polymerase chain reaction indicates a trimodular genotype-phenotype correlation. *Pharmacogenetics* 10: 557-565.
6. Pastore, A., Piemonte, F., Locatelli, M., Lo Russo, A., Gaeta, L.M., Tozzi, G. and Federici, G. 2001. Determination of blood total, reduced, and oxidized glutathione in pediatric subjects. *Clin. Chem.* 47: 1467-1469.
7. Pompella, A., Visvikis, A., Paolicchi, A., De Tata, V. and Casini, A.F. 2003. The changing faces of glutathione, a cellular protagonist. *Biochem. Pharmacol.* 66: 1499-1503.

CHROMOSOMAL LOCATION

Genetic locus: GSTT2 (human) mapping to 22q11.23.

PRODUCT

GSTT2 (h3): 293T Lysate represents a lysate of human GSTT2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

GSTT2 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive GSTT2 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.