



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

POU3F4 (h2): 293T Lysate: sc-129522

BACKGROUND

The Brn family of transcription factors are found in a highly restricted subset of neurons and are critical to the early embryonic development of the central nervous system. POU3F4 (POU domain, class 3, transcription factor 4), also known as OTF9, DFN3, DFNX2, BRAIN-4 or Brn-4, is a 361 amino acid class III POU domain protein that belongs to the Brn family. Localized to the nucleus, POU3F4 contains one POU-specific domain and one homeobox DNA-binding domain through which it is thought to function as a brain-specific transcription factor that effects neuronal development. Defects in the gene encoding POU3F4 are associated with X-linked deafness type 3 (DFN3), a disorder characterized by both progressive sensorineural deafness and conductive hearing loss caused by stapes gushers (large vestibular aqueducts).

REFERENCES

1. Malik, K.F., Jaffe, H., Brady, J. and Young, W.S. 1997. The class III POU factor Brn-4 interacts with other class III POU factors and the heterogeneous nuclear ribonucleoprotein U. *Brain Res. Mol. Brain Res.* 45: 99-107.
2. de Kok, Y.J., Cremers, C.W., Ropers, H.H. and Cremers, F.P. 1997. The molecular basis of X-linked deafness type 3 (DFN3) in two sporadic cases: identification of a somatic mosaicism for a POU3F4 missense mutation. *Hum. Mutat.* 10: 207-211.
3. Phippard, D., Heydemann, A., Lechner, M., Lu, L., Lee, D., Kyin, T. and Crenshaw, E.B. 1998. Changes in the subcellular localization of the Brn4 gene product precede mesenchymal remodeling of the otic capsule. *Hear. Res.* 120: 77-85.
4. Hagiwara, H., Tamagawa, Y., Kitamura, K. and Kodera, K. 1998. A new mutation in the POU3F4 gene in a Japanese family with X-linked mixed deafness (DFN3). *Laryngoscope* 108: 1544-1547.
5. Shimazaki, T., Arsenijevic, Y., Ryan, A.K., Rosenfeld, M.G. and Weiss, S. 1999. A role for the POU-III transcription factor Brn-4 in the regulation of striatal neuron precursor differentiation. *EMBO J.* 18: 444-456.
6. Xia, A.P., Kikuchi, T., Minowa, O., Katori, Y., Oshima, T., Noda, T. and Ikeda, K. 2002. Late-onset hearing loss in a mouse model of DFN3 non-syndromic deafness: morphologic and immunohistochemical analyses. *Hear. Res.* 166: 150-158.

CHROMOSOMAL LOCATION

Genetic locus: POU3F4 (human) mapping to Xq21.1.

PRODUCT

POU3F4 (h2): 293T Lysate represents a lysate of human POU3F4 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

POU3F4 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive POU3F4 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.