



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TFIP11 (h): 293T Lysate: sc-129936

BACKGROUND

TFIP11 (tuftelin-interacting protein 11), also known as NTR1, TIP39, STIP or hNtr1, is an 837 amino acid protein that belongs to the TFP11/STIP family. TFIP11 localizes to the nucleus as well as the cytoplasm and contains one G-patch domain, which is suggested to be a highly conserved domain of many RNA-processing proteins. Considered a novel splicing factor, TFIP11 may be involved in spliceosome disassembly and may act as a subnuclear storage compartment for splicing components. As a possible enamel protein, TFIP11 is thought to play a role in the differentiation of ameloblasts and odontoblasts or in the formation of the enamel extracellular matrix. Two isoforms of TFIP11 exist due to alternative splicing events.

REFERENCES

1. Usdin, T.B., Wang, T., Hoare, S.R., Mezey, E. and Palkovits, M. 2000. New members of the parathyroid hormone/parathyroid hormone receptor family: the parathyroid hormone 2 receptor and tuberoinfundibular peptide of 39 residues. *Front. Neuroendocrinol.* 21: 349-383.
2. Paine, C.T., Paine, M.L., Luo, W., Okamoto, C.T., Lyngstadaas, S.P. and Snead, M.L. 2000. A Tuftelin-interacting protein (TIP39) localizes to the apical secretory pole of mouse ameloblasts. *J. Biol. Chem.* 275: 22284-22292.
3. Usdin, T.B., Bonner, T.I. and Hoare, S.R. 2002. The parathyroid hormone 2 (PTH2) receptor. *Recept. Channels* 8: 211-218.
4. Wen, X., Lei, Y.P., Zhou, Y.L., Okamoto, C.T., Snead, M.L. and Paine, M.L. 2005. Structural organization and cellular localization of Tuftelin-interacting protein 11 (TFIP11). *Cell. Mol. Life Sci.* 62: 1038-1046.
5. Wang, H.J., Tannukit, S., Wen, X., Shapiro, J.L., Snead, M.L. and Paine, M.L. 2006. Using the yeast two-hybrid assay to discover protein partners for the leucine-rich Amelogenin peptide and for Tuftelin-interacting protein 11. *Eur. J. Oral Sci.* 114: 276-279.
6. Herrmann, G., Kais, S., Hoffbauer, J., Shah-Hosseini, K., Brüggelolte, N., Schober, H., Fäsi, M. and Schär, P. 2007. Conserved interactions of the splicing factor NTR1/Spp382 with proteins involved in DNA double-strand break repair and telomere metabolism. *Nucleic Acids Res.* 35: 2321-2332.
7. Patir, A., Seymen, F., Yildirim, M., Deeley, K., Cooper, M.E., Marazita, M.L. and Vieira, A.R. 2008. Enamel formation genes are associated with high caries experience in Turkish children. *Caries Res.* 42: 394-400.
8. Tannukit, S., Wen, X., Wang, H. and Paine, M.L. 2008. TFIP11, CCNL1 and EWSR1 protein-protein Interactions, and their nuclear localization. *Int. J. Mol. Sci.* 9: 1504-1514.
9. Wen, X., Tannukit, S. and Paine, M.L. 2008. TFIP11 Interacts with mDEAH9, an RNA Helicase Involved in spliceosome disassembly. *Int. J. Mol. Sci.* 9: 2105-2113.

CHROMOSOMAL LOCATION

Genetic locus: TFIP11 (human) mapping to 22q12.1.

PRODUCT

TFIP11 (h): 293T Lysate represents a lysate of human TFIP11 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

TFIP11 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive TFIP11 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.