



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ZFP820 siRNA (m): sc-155590

BACKGROUND

Santa Cruz Biotechnology now offers siRNA and shRNA (plasmid and lentiviral) gene silencers targeted to all (100%) known human and mouse genes. siRNAs are provided as pools of 3 target specific 19-25 nt siRNAs designed to knock down expression of specific gene of interest. Each vial contains 3.3 nmol of lyophilized siRNA and is sufficient for 50-100 transfections. For independent verification of gene silencing results, we also will provide, upon request, individual siRNA duplex components, each containing 3.3 nmol of lyophilized siRNA. shRNA Plasmid DNA Gene Silencers are provided as pools of three to five lentiviral vector plasmids each encoding target-specific 19-25 nt (plus hairpin) shRNA (sequences correspond to our well-established existing product line of siRNA gene silencers). shRNA Plasmids are provided as 20 µg transfection-ready plasmid DNA for either transient or stable (puromycin-selection) target specific gene silencing. shRNA Lentiviral Particles are provided as pools of three to five expression constructs each encoding target-specific 19-25 nt (plus hairpin loop) shRNA and are provided as 200 µl transduction-ready viral particles for gene silencing in mammalian cells (human or mouse). Suitable control antibodies and RT-PCR primers are available for monitoring gene expression knockdown.

CHROMOSOMAL LOCATION

Genetic locus: Zfp820 (mouse) mapping to 17 A3.3.

PRODUCT

ZFP820 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZFP820 shRNA Plasmid (m): sc-155590-SH and ZFP820 shRNA (m) Lentiviral Particles: sc-155590-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

ZFP820 siRNA (m) is recommended for the inhibition of ZFP820 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZFP820 gene expression knockdown using RT-PCR Primer: ZFP820 (m)-PR: sc-155590-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.