



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

caspase-3 (h4): 293 Lysate: sc-158344

BACKGROUND

Caspase-3, also known as apopain, SCA-1, Yama and CPP32, is an aspartate-specific cysteine protease that belongs to the ICE subfamily of caspases. Caspase-3 is expressed in cells as an inactive precursor from which the p17 and p11 subunits of the mature caspase-3 are proteolytically generated during apoptosis. The caspase-3 precursor is first cleaved at Asp 175-Ser 176 to produce the p11 subunit and the p20 peptide. Subsequently, the p20 peptide is cleaved at Asp 28-Ser 29 to generate the mature p17 subunit. The active caspase-3 enzyme is a heterodimer composed of two p17 and two p11 subunits. At the onset of apoptosis, caspase-3 proteolytically cleaves PARP at a Asp 216-Gly 217 bond. During the execution of the apoptotic cascade, activated caspase-3 releases SREBP from the membrane of the ER in a proteolytic reaction that is distinct from their normal sterol-dependent activation. Caspase-3 cleaves and activates SREBPs between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase-3 also cleaves and activates caspase-6, -7 and -9. The human caspase-3 gene encodes a cytoplasmic protein that is highly expressed in lung, spleen, heart, liver, kidney and cells of the immune system.

REFERENCES

- Nicholson, D., et al. 1995. Identification and inhibition of the ICE/CED-3 protease necessary for mammalian apoptosis. *Nature* 37: 37-43.
- Cohen, G.M. 1997. Caspases: the executioners of apoptosis. *Biochem. J.* 326: 1-16.
- Higgins, M.E., et al. 2001. Apoptosis-induced release of mature sterol regulatory element-binding proteins activates sterol-responsive genes. *J. Lipid Res.* 42: 1939-1946.
- Vallender, E.J. and Lahn, B.T. 2006. A primate-specific acceleration in the evolution of the caspase-dependent apoptosis pathway. *Hum. Mol. Genet.* 15: 3034-3040.
- Liu, X., et al. 2006 Apoptosis and proliferation markers in diffusely infiltrating astrocytomas: profiling of 17 molecules. *J. Neuropathol. Exp. Neurol.* 65: 905-913.
- Floros, K.V., et al. 2006. Alterations in mRNA expression of apoptosis-related genes BCL2, BAX, FAS, caspase-3, and the novel member BCL2L12 after treatment of human leukemic cell line HL60 with the antineoplastic agent etoposide. *Ann. N.Y. Acad. Sci.* 1090: 89-97.
- Kriebel, F., et al. 2006. Caspase-3 activation, Bcl-2 contents, and soluble FAS-ligand are not related to the inflammatory marker profile in patients with sepsis and septic shock. *Ann. N.Y. Acad. Sci.* 1090: 168-176.
- Schneider, E.M., et al. 2006. Role of ATP in trauma-associated cytokine release and apoptosis by P2X7 ion channel stimulation. *Ann. N.Y. Acad. Sci.* 1090: 245-252.
- Das, U., et al. 2007. 3-(3,4,5-trimethoxyphenyl)-1-oxo-2-propene: A novel pharmacophore displaying potent multidrug resistance reversal and selective cytotoxicity. *Bioorg. Med. Chem.* 15: 3373-3380.

CHROMOSOMAL LOCATION

Genetic locus: CASP3 (human) mapping to 4q35.1.

PRODUCT

caspase-3 (h4): 293 Lysate represents a lysate of human caspase-3 transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

caspase-3 (h4): 293 Lysate is suitable as a Western Blotting positive control for human reactive caspase-3 antibodies. Recommended use: 10-20 µl per lane.


Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

caspase-3 (3F49): sc-70498 is recommended as a positive control antibody for Western Blot analysis of enhanced human caspase-3 expression in caspase-3 transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



caspase-3 (3F49): sc-70498. Western blot analysis of procaspase-3 expression in non-transfected: sc-110760 (A) and human caspase-3 transfected: sc-158344 (B) 293 whole cell lysates.



caspase-3 (3CSP01): sc-65496. Western blot analysis of procaspase-3 expression in non-transfected: sc-110760 (A) and human caspase-3 transfected: sc-158344 (B) 293 whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.