



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cdk4 (h3): 293 Lysate: sc-158369

BACKGROUND

Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Several Cdk proteins have been identified, including Cdk2-Cdk8, PCTAIRE-1-PCTAIRE-3, PITALRE and PITSLRE. Cdk4, in complex with D-type cyclins, is thought to regulate cell growth during the G₁ phase of the cell cycle. This association with a D-type cyclin upregulates Cdk4 activity, whereas binding to the Cdk inhibitor p16 downregulates Cdk4 activity. Activation of the Cdk4-cyclin complexes requires phosphorylation on a single threonyl residue of Cdk4, catalyzed by a Cdk-activating protein (CAK).

REFERENCES

- Okuda, T., et al. 1992. PCTAIRE-1 and PCTAIRE-2: two members of a novel Cdc2/Cdc28-related protein kinase gene family. *Oncogene* 7: 2249-2258.
- Serrano, M., et al. 1993. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/Cdk4. *Nature* 366: 704-707.
- Pines, J. 1994. The cell cycle kinases. *Semin. Cancer Biol.* 5: 305-313.
- Kato, J.Y., et al. 1994. Regulation of cyclin D-dependent kinase (Cdk4) by Cdk4-activating kinase. *Mol. Cell. Biol.* 14: 2713-2721.
- Matsuoka, M., et al. 1994. Activation of cyclin-dependent kinase 4 (Cdk4) by mouse MO15-associated kinase. *Mol. Cell. Biol.* 14: 7265-7275.
- MacLachlan, T.K., et al. 1995. Cyclins, cyclin-dependent kinases and Cdk inhibitors: implications in cell cycle control and cancer. *Crit. Rev. Eukaryot. Gene Expr.* 5: 127-156.
- Siebert, R., et al. 1996. Role of the cyclin-dependent kinase 4 and 6 inhibitor gene family p15, p16, p18 and p19 in leukemia and lymphoma. *Leuk. Lymphoma* 23: 505-520.
- Dirks, P.B., et al. 1997. Current concepts in neuro-oncology: the cell cycle—a review. *Neurosurgery* 40: 1000-1013.
- Coleman, K.G., et al. 1997. Identification of Cdk4 sequences involved in cyclin D1 and p16 binding. *J. Biol. Chem.* 272: 18869-18874.

CHROMOSOMAL LOCATION

Genetic locus: CDK4 (human) mapping to 12q14.1.

PRODUCT

Cdk4 (h3): 293 Lysate represents a lysate of human Cdk4 transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Cdk4 (h3): 293 Lysate is suitable as a Western Blotting positive control for human reactive Cdk4 antibodies. Recommended use: 10-20 µl per lane.


Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

Cdk4 (DCS-156): sc-53636 is recommended as a positive control antibody for Western Blot analysis of enhanced human Cdk4 expression in Cdk4 transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Cdk4 (DCS-156): sc-53636. Western blot analysis of Cdk4 expression in non-transfected: sc-110760 (A) and human Cdk4 transfected: sc-158369 (B) 293 whole cell lysates.



Cdk4 (3F121): sc-70831. Western blot analysis of Cdk4 expression in non-transfected: sc-110760 (A) and human Cdk4 transfected: sc-158369 (B) 293 whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.