

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

granzyme B (h3): 293T Lysate: sc-158574



BACKGROUND

Granzyme A and granzyme B are serine proteases that mediate apoptotic signaling in cytotoxic T lymphocytes (CTL) and natural killer (NK) cells. Both granzyme A and granzyme B are synthesized as inactive proenzymes, and they are stored within cytolytic granules and released by effector cells during degranulation. In activated CTLs, granzyme A and granzyme B are processed and activated by cathepsin C, and they then function to induce apoptosis by two distinct pathways. Granzyme B proteolytically cleaves and activates members of the caspase family of cysteine proteases, including caspase-3, caspase-6, caspase-7 and caspase-9. When cleaved, these caspases assemble into active holoenzymes that then mediate apoptosis through a defined proteolytic cascade involving nuclear lamins and PARP (poly (ADP ribose) polymerase). Granzyme A mediates the activation of apoptosis by inducing single-strand DNA breaks, membrane perturbation and nuclear condensations in an alternative pathway that is independent from caspase activation or the caspase proteolytic cascade.

REFERENCES

- Gershenfeld, H.K., et al. 1988. Cloning and chromosomal assignment of a human cDNA encoding a T cell- and natural killer cell-specific trypsin-like serine protease. Proc. Natl. Acad. Sci. USA 85: 1184-1188.
- Shresta, S., et al. 1995. Natural killer and lymphokine-activated killer cells require granzyme B for the rapid induction of apoptosis in susceptible target cells. Proc. Natl. Acad. Sci. USA 92: 5679-5683.
- Trapani, J.A., et al. 1996. A putative role in the mechanism of cytotoxic lymphocyte-mediated apoptosis. Localization of granzyme B in the nucleus. J. Biol. Chem. 271: 4127-4133.
- Atkinson, E.A., et al. 1998. Cytotoxic T lymphocyte-assisted suicide. Caspase-3 activation is primarily the result of the direct action of granzyme B. J. Biol. Chem. 273: 21261-21266.
- Trapani, J.A., et al. 1998. Efficient nuclear targeting of granzyme B and the nuclear consequences of apoptosis induced by granzyme B and perforin are caspase-dependent, but cell death is caspase-independent. J. Biol. Chem. 273: 27934-27938.
- Pham, C.T., et al. 1999. Dipeptidyl peptidase I is required for the processing and activation of granzymes A and B *in vivo*. Proc. Natl. Acad. Sci. USA 96: 8627-8632.
- 7. Shresta, S., et al. 1999. Granzyme A initiates an alternative pathway for granule-mediated apoptosis. Immunity 10: 595-605.

CHROMOSOMAL LOCATION

Genetic locus: GZMB (human) mapping to 14q12.

PRODUCT

granzyme B (h3): 293T Lysate represents a lysate of human granzyme B transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

granzyme B (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive granzyme B antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

granzyme B (GRB7): sc-73620 is recommended as a positive control antibody for Western Blot analysis of enhanced human granzyme B expression in granzyme B transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA





granzyme B (GRB7): sc-73620. Western blot analysis of granzyme B expression in non-transfected: sc-117752 (**A**) and human granzyme B transfected: sc-158574 (**B**) 293T whole cell lysates. granzyme B (GB7): sc-56119. Western blot analysis of granzyme B expression in non-transfected: sc-117752 (**A**) and human granzyme B transfected: sc-158574 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.